The B. H. Hill Library



North Carolina State University

QK190

F6

THIS BOOK IS DUE ON THE DATE INDICATED BELOW AND IS SUBJECT TO AN OVERDUE FINE AS POSTED AT THE CIRCULATION DESK.







FLORA OF BURLINGTON AND VICINITY


A list of the

FERN AND SEED PLANTS

Growing without cultivation

Nellie F. Flynn

BURLINGTON:
Free Press Printing Company,
1911



My Husband



FLORA OF BURLINGTON AND VICINITY

PREFACE

This list was begun at the suggestion of Dr. L. R. Jones, formerly of the University of Vermont, now of the University of Wisconsin. It was intended to be accompanied by an account of the plant geography and ecology of the region by Dr. Clifton D. Howe. Since the removal of Dr. Howe to Toronto has delayed the completion of his surveys* it seems best to publish this now as a preliminary list. It is hoped that when Dr. Howe's report is completed, a revision of this list may be published with it.

The list is intended to include the names of all the fern and seed plants which are natives of or grow without cultivation in Burlington and its vicinity, including with Burlington, the towns of South Burlington, Colchester, Essex, Williston and Shelburne. Each name has an extant specimen back of it, except in three or four cases which are specifically mentioned. In these cases we are sure of the identity of the plant cited. Except in a few instances, specimens are in the herbarium of the University of Vermont or in the private collection of the author.

To stimulate further search, a supplementary list has been added containing the names of additional plants which have been doubtfully reported for the region or which for reasons named should be sought here. In the case of rare plants, stations and names of collectors have been given. For the sake of brevity, initials of collectors' names have not been given except where there are two of like name. In order to complete the records, the full name, and present address of the living collectors where known, are here given.

^{*}Dr. Howe has recently published certain of his observations on the plant geography of this region entitled: The Reforestation of the Sand Plains of Vermont. Botanical Gazette, Feb'y, 1910.

Rev. J. A. Bates, South Royalston, Mass.; Hon. R. D. Benedict, Burlington; Lucius Bigelow, Burlington; William Boott (specimens in Gray Herbarium); Dr. Ezra Brainerd, Middlebury, Vt.; Miss Martha Brookes-Brown (specimens in U. V. M. Herbarium); C. E. Bryant, Williston, Vt.; Dr. G. P. Burns, University of Vermont, Burlington; H. B. Chittenden (specimens in U. V. M. Herbarium); H. J. Collins, West Lebanon, N. H.; A. W. Edson (specimens in U. V. M. Herbarium); W. W. Eggleston, Washington, D. C.; Mrs. N. F. Flynn, Burlington; Mrs. G. I. Forbes, Burlington; Mrs. E. F. Gebhardt, Shelburne, Vt.: C. M. Gifford, Morgantown, W. Va.; A. H. Gilbert, Washington, D. C.; Miss M. K. Goddard, Lexington, Mass.; D. B. Griffin, Winooski, Vt.; Dr. A. J. Grout, New Dorp, N. Y.; Mrs. A. J. Grout, New Dorp, N. Y.; Dr. T. E. Hazen, Barnard College, New York, N. Y.; J. P. Helyer, University of Vermont, Burlington; C. G. Hinsdale, Boston, Mass.; Miss Miriam Hitchcock, Pittsford, Vt.; F. M. Hollister, Tewksbury, Mass.; Mrs. Henry Holt, Burlington; S. C. Hood, Orange City, Fla.: F. H. Horsford, Charlotte, Vt.; Dr. C. D. Howe, Toronto University, Toronto, Can.; F. E. Hubbard, Keene Valley, N. Y.; Dr. L. R. Jones, University of Wisconsin, Madison: G. L. Kirk, Rutland, Vt.; W. F. Macrae (specimens in Gray Herbarium); Miss Lizzie Moore, Burlington; Prof. W. J. Morse, University of Maine, Orono, Me.; William Oakes (specimens in Gray Herbarium); W. A. Orton, Washington, D. C.; Prof. G. H. Perkins, University of Vermont, Burlington: H. L. Priest (specimens in U. V. M. Herbarium); Dr. C. G. Pringle, University of Vermont, Burlington; Mrs. F. A. Richardson, Burlington; Dr. J. W. Robbins (specimens in Gray Herbarium); F. A. Ross (specimens in U. V. M. Herbarium); J. K. Shaw, Amherst, Mass.; Dr. L. P. Sprague, Chateaugay, N. Y.; R. R. Strait, Franklin, Vt.; Prof. H. A. P. Torrey (specimens in U. V. M. Herbarium); Prof. Joseph Torrey (specimens in U. V. M. Herbarium); Miss Phoebe Towle, Burlington; C. C. Tracy (specimens in U. V. M. Herbarium); Mrs. F. H. Walker, Portland, Ore.; Rev. Levi Wild, Royalton, Vt.; Mrs. W. H. Zottman, Burlington.

It is a matter of regret, that the older botanists who were the first collectors of this region, namely, Macrae, Robbins, Torrey and Benedict, who collected here about the middle of the last century, have so few authenticated specimens in herbaria, that due credit cannot be given them. It is hoped that anyone having such specimens or those of other collectors which are not fully credited, will inform the author.

The seventh edition of Gray's Manual has been followed strictly in nomenclature, except where for the sake of brevity, initials have been eliminated or names abbreviated, and synonyms have only been given where corrections have been made since the Manual came out. Common names have been inserted freely as befitting the general use which may be made of such a list.

Following the usage of the Vermont Flora, four adjectives, namely, "common," "frequent," "occasional" and "rare," have been adopted to denote frequency of occurrence. These are intended to be associated with the habitat—thus, "sandy soil, common," means, common on sandy soils, though it may be rare in other habitats. The term "rare" is used for those plants which occur in but few stations and there but sparingly. When they are abundant in such stations, the word "local" has been preferred. The habitats given are based on personal observation where possible.

Following the common practice, small capitals have been used for names of plants not native to the region as far as our knowledge goes. In a few cases these may be native of some other part of the state, but introduced here.

This list is necessarily provisional and incomplete, and will have served its best purpose if it prepares the way for a fuller one later. To that end it is hoped that anyone who can give added information about the plants in this list, or who finds any additional species, will let the author know, and deposit specimens either with her or in the herbarium of the University of Vermont. Proper recognition will be made of such assistance in the proposed revision.

My thanks are due not only to Dr. L. R. Jones who has been of the greatest help and encouragement, but to Dr. Ezra Brainerd, Mr. W. W. Eggleston, Rev. J. A. Bates, Dr. T. E. Hazen and many others for assistance given. Thanks are also due to Prof. M. L. Fernald, who has examined and verified many doubtful specimens.

NELLIE F. FLYNN.



CONTENTS

	Page
Catalogue of Flowering Plants and Ferns	. 1
Pteridophytes	. 1
Spermatophytes	. 6
Supplementary List	. 94
Statistical Summaries	. 101
Additions and Corrections	103
Index	104



Flora of Burlington and Vicinity

PTERIDOPHYTA Ferns and Fern Allies

POLYPODIACEAE. FERN FAMILY

ADIANTUM (Tourn.) L. MAIDENHAIR

- A. pedatum L. Rich moist woods; frequent. July.

 ASPIDIUM Sw. SHIELD FERN. WOOD FERN.
- A. Boottii Tuck. Boott's Shield Fern. Wet thickets and about ponds; rare. July. Burlington, (N. F. F.); Essex and Colchester, (Hazen).
- A. cristatum (L.) Sw. Crested Shield Fern. Moist thickets and swamps; occasional. July.
- A. cristatum (L.) Sw., var. Clintonianum Eaton. Clinton's Shield Fern. Swampy woods; rare. July. Colchester, (N. F. F.).
- A. Goldianum Hook. Goldie's Fern. Rich woods; rare. July. Shelburne.
- A. marginale (L.) Sw. Marginal Shield Fern, Evergreen Wood Fern. Rocky hillsides in rich woods; common. July, Aug.
- A. noveboraceuse (L.) Sw. New York Fern. Rich woods; common. Aug.
- A. spinulosum (Müller) Sw. Spinulose Shield Fern. Rich woods; occasional, July.
- A. spinulosum (Müller) Sw., var. intermedium (Muhl.) Eaton. Woods; common. July.
- A. Thelypteris (L.) Sw. Marsh Fern. Marshes: common. Aug.

ASPLENIUM L. SPLEENWORT

- A. acrostichoides Sw. Silvery Spleenwort. Rich woods; rare. July, Aug. Shelburne, (Bates); Burlington, (N. F. F.).
- A. angustifolium Michx. Narrow-leaved Spleenwort. Rich woods; rare. Aug. Shelburne.
- A. Filix-femina (L.) Bernh. Lady Fern. Moist woods; common in several different forms. July, Aug.
- A. platyneuron (L.) Oakes. Ebony Spleenwort. Rocky open woods; occasional. July, Aug. A form near var. incisum (Howe) Robins. occurs, Colchester, (Griffin).
- A. Ruta-muraria L. Rue Spleenwort. Limestone cliffs; occasional.

 July.
- A. Trichomanes L. Maidenhair Spleenwort. Shaded rocks; occasional. July. A form near var. INCISUM Moore occurs, Colchester, (Griffin).

CAMPTOSORUS Link. WALKING LEAF

('. rhizophyllus (L.) Link. Shaded rocks, especially limestone; occasional. Aug. Forms occur with the fronds forked at the tip and with the auricles greatly elongated.

CRYPTOGRAMMA R. Br. ROCK BRAKE

C. Stelleri (Gmel.) Prantl. Slender Rock Brake. Moist limestone rocks; rare. July. South Burlington, (Perkins).

CYSTOPTERIS Bernh. BLADDER FERN

- C. bulbifera (L.) Bernh, Common Bladder Fern. Moist shaded ravines; frequent. July.
- C. fragilis (L.) Bernh. Fragile Bladder Fern. Rocky woods and ravines; frequent. June.

DICKSONIA L'Hér.

D. punctilobula (Michx.) Gray. Hay-scented Fern. Moist shaded places; occasional. Aug.

ONOCLEA L.

- 0. sensibilis L. Sensitive Fern. Moist meadows and thickets; common. Sterile June; fertile Sept. Sports are occasional,—the so-called var. OBTUSILOBATA (Schkuhr) Torr. Burlington, (Miss Goddard); South Burlington, (Ross).
- 0. Struthiopteris (L.) Hoffm. Ostrich Fern. Alluvial soil; common. Sterile June; fertile Sept.

PELLAEA Link. CLIFF BRAKE

P. atropurpurea (L.) Link. Purple Cliff Brake. Limestone cliffs; occasional. July.

PHEGOPTERIS (Presl) Fée. BEECH FERN

- P. Dryopteris (L.) Fée. Oak Fern. Moist woods; frequent. July.
- P. hexagonoptera (Michx.) Fée. Broad Beech Fern. Rich woods; rare. July, Aug. Shelburne, (Jones); Burlington, (N. F. F.); Colchester, (Griffin).
- P. polypodioides Fée. Long Beech Fern. Damp woods; occasional. July.

POLYPODIUM (Tourn.) L. POLYPODY

I'. vulgare L. Rocks; common. July, Aug.

POLYSTICHUM Roth

P. acrostichoides (Michx.) Schott. Christmas Fern. Rocky woods; common. July. Var. Schweinitzh (Beck) Small. (Var. incisum Gray) Occasional. A crisped form occurs, Burlington, (Mrs. Richardson, Ross); Colchester, (Griffin).

PTERIS L. BRAKE OF BRACKEN

P. aquilina L. Pastures and woodlands; common, Aug. Var. PSEUDO-CAUDATA Clute. Rare. Colchester, (Ross).

WOODSIA R. Br.

- W. ilvensis (L.) R. Br. Rusty Woodsia. Exposed rocks; occasional. July. Abundant at one station.
- W. obtusa (Spreng.) Torr. Blunt-lobed Woodsia. Rocky banks; occasional. July.

WOODWARDIA Sm. CHAIN FERN

W. virginica (L.) Sm. Virginia Chain Fern. Wet swamp, Colchester. (Robbins). July.

OSMUNDACEAE. FLOWERING FERN FAMILY

OSMUNDA (Tourn.) L. FLOWERING FERN

O. cinnamomea L. Cinnamon Fern. Wet woods and pastures; common. Fertile May; sterile June. Variable in the shape and cutting of the pinnules. Var. INCISA Hunt. Rare. Burlington, (Ross, N. F. F.); Colchester, (Griffin).

- Claytoniana L. Interrupted Fern. Wet woods and pastures; common. Fertile May; sterile June.
- (), regalis L. Royal Fern. Wet woods and swamps; common. June. Forms occur with the sterile fronds fertile at the tips and with the pinnules incised; South Burlington and Burlington, (N. F. F.).

OPHIOGLOSSACEAE. ADDER'S TONGUE FAMILY

BOTRYCHIUM Sw. MOONWORT

- B. lanceolatum (Gmel.) Angst., var. augustisegmentum Pease & Moore. Lance-leaved Grape Fern. Cool open woods; rare. July. Burlington, (Ross).
- B. obliquum Muhl. Grape Fern. Moist sandy soil; occasional. Sept. Var. DISSECTUM (Spreng.) Clute. Occasional. Sept.
- B. ramosum (Roth) Aschers. Rich woods; rare. July. Burlington, (Ross); Colchester, (Griffin).
- B. simplex Hitchc. Little Grape Fern. Moist pastures; rare. June, July. Burlington, (N. F. F., Ross).
- B. ternatum (Thunb.) Sw., var. intermedium Eaton. Ternate Grape Fern. Sandy soil; rare. Sept. Burlington, (N. F. F., Ross). Var. RUTAEFOLIUM (A. Br.) Eaton. Rare. Sept. Burlington, (Gilbert).
- B. virginianum (L.) Sw. Rattlesnake Fern. Rich woods; frequent and variable in size. June, July.

OPHIOGLOSSUM (Tourn.) L. ADDER'S TONGUE

vulgatum L. Moist pastures; rare. June, July. Burlington, (N. F. F.); Colchester, (Ross).

EQUISETACEAE. HORSETAIL FAMILY

, EQUISETUM (Tourn.) L. HORSETAIL

- E. arvense L. Common Horsetail. Moist gravelly soil; common. Fertile Apr., May; sterile June.
- E. fluviatile L. Pipes. Shallow water with muddy bottom; common. Fertile June; sterile July. A form occurs the branches of which bear numerous small spikes; Burlington, (N. F. F.).
- E. hyemale L., var. affine (Engelm.) Eaton. Common Scouring Rush. Moist gravelly banks; common. May-Sept.

- E. hyemale L., var. intermedium Eaton. Scouring Rush. Moist sandy soil; rare. May-Aug. Burlington, (Howe).
- E. litorale Kühl. Shore Horsetail. Gravelly shores of Lake Champlain; rare. May, June. Burlington, (Grout and Jones).
- E. palustre L. Marsh Horsetail. Wet sandy shore of Lake Champlain; rare. June, July. Burlington, (Grout and Jones).
- E. scirpoides Michx. Cool moist woods; occasional. May, June.
- E. sylvaticum L. Wood Horsetail. Moist shady places; frequent. Fertile May; sterile June.
- E. variegatum Schleich. Gravelly shores of Lake Champlain; rare. May, June. Burlington, (Eggleston and Grout, N. F. F.); Colchester, (Sprague).

LYCOPODIACEAE. CLUB MOSS FAMILY

LYCOPODIUM (Rupp.) L. CLUB Moss

- L. annotinum L. Stiff Club Moss. Cold woods; rare. July, Aug. Burlington, (Grout).
- I. clavatum L. Common Club Moss. Dry woods; common. Aug., Sept.
- L. complanatum L., var. flabelliforme Fern. Ground Pine. Dry woods; frequent. Sept.
- L. inundatum L. Bog Club Moss. In sphagnum; rare. Aug., Sept. Williston, (Pringle); South Burlington, (Ross); Colchester and Burlington, (N. F. F.).
- L. lucidulum Michx. Shining Club Moss. Cold damp woods; frequent. Sept.
- L. obscurum L. Tree Club Moss. Moist woods; rare. Sept. South Burlington, (Hood).
- L. obscurum L., var. dendroideum (Michx.) Eaton. Common Tree Club Moss. Moist woods; common. Sept.
- L. tristachyum Pursh. Ground Pine. Dry sandy soil; rare. July, Aug. Colchester and Essex, (Hazen).

SELAGINELLACEAE

SELAGINELLA Beauv.

S. rupestris (L.) Spring. Rock Selaginella. Dry exposed rocks; occasional. Aug., Sept.

SPERMATOPHYTA

Seed-Plants, Phanerogamia, or Flowering Plants

TAXACEAE. YEW FAMILY

TAXUS (Tourn.) L. YEW

T. canadensis Marsh. American Yew, Ground Hemlock. Moist banks; frequent. Apr., May.

PINACEAE. PINE FAMILY

ABIES (Tourn.) Hill. FIR

A. balsamea (L.) Mill. Balsam or Balm-of-Gilead Fir. Damp woods; rare. May. Burlington and South Burlington, (Hubbard); Essex, (Jones).

JUNIPERUS (Tourn.) L. JUNIPER

- J. communis L., var. depressa Pursh. Common Juniper. Dry sterile hills; frequent. Apr., May.
- J. virginiana L. Red Cedar or Savin. Dry rocky hills; frequent. May.

LARIX (Tourn.) Mill. LARCH

L. larieina (Du Roi) Koch. American or Black Larch, Tamarack, Hackmatack. Cold swamps; common. May.

PICEA Link. SPRUCE

- P. canadensis (Mill.) BSP. White or Cat Spruce. A few trees at one station, Colchester, (Griffin); Essex, (Howe). May.
- P. mariana (Mill.) BSP. Bog or Black Spruce. Sphagnum swamps; common. May.

P. rubra (Du Roi) Dietr. Red Spruce. Frequent in deep ravines about Burlington, (Howe), but rare elsewhere. May. South Burlington, (Hubbard).

PINUS (Tourn.) L. PINE

- P. resinosa Ait. Red or Norway Pine. Dry woods; frequent. May, June.
- P. rigida Mill. Pitch Pine. Barren sandy soil; common. May, June.
- P. Strobus L. White Pine, Common, June.

THUJA L. ARBOR VITAE

T. occidentalis L. Arbor Vitae, White Cedar. Swamps and rocky banks; common. Apr., May.

TSUGA (Endl.) Carr. HEMLOCK

T. canadensis (L.) Carr. Swamps and rocky woods; common. May.

TYPHACEAE. CAT-TAIL FAMILY

TYPHA (Tourn.) L. CAT-TAIL FLAG

- T. angustifolia L. Narrow-leaved Cat-tail. Marshes; rare. July. Burlington, (N. F. F.); Colchester, (Griffin).
- T. latifolia L. Common Cat-tail. Marshes; common. July.

SPARGANIACEAE. BUR=REED FAMILY

SPARGANIUM (Tourn.) L. BUR-REED

- S. americanum Nutt. Muddy shores; rare. July, Aug. Burlington, (N. F. F.).
- S. diversifolium Graeb. Muddy shores; rare. July, Aug. Colchester, (N. F. F.).
- S. eurycarpum Engelm. Broad-fruited Bur-reed. Borders of ponds and rivers; frequent. July, Aug.

NAJADACEAE. PONDWEED FAMILY

NAJAS L. NAIAD

N. flexilis (Willd.) Rostk. & Schmidt. Ponds and slow streams; common. July-Sept.

POTAMOGETON (Tourn.) L. PONDWEED

- P. americanus C. & S. Mouth of Winooski River, (Grout and Tracy).
 July, Aug.
- P. amplifolius Tuck. Large-leaved Pondweed. Shelburne Pond, (Jones); Colchester, (Griffin). July, Aug.
- P. CRISPUS L. Malletts Bay, (Griffin). July, Aug.
- P. dimorphus Raf. Ponds and slow streams; frequent. July, Aug.
- P. epihydrus Raf. Still or flowing water; frequent. July, Aug.
- P. foliosus Raf. Still or flowing water; rare. July, Aug. Burlington and Colchester, (N. F. F.).
- P. Friesii Rupr. Shelburne Pond, (Pringle). Aug., Sept.
- H heterophyllus Schreb. Mouth of Winooski River, (Grout and Tracy, Howe); Malletts Bay, (N. F. F.). July, Aug. Forma TERRESTRIS Schlecht. Exsiccated place, Burlington, (N. F. F.).
- I'. natans L. Common Pondweed. Shelburne Pond, (Jones); Malletts Bay, (Griffin). July, Aug.
- P. pectinatus L. Malletts Bay, (Griffin). July-Sept.
- P. perfoliatus L. Slow streams and shallow waters of Lake Champlain; frequent. Sept.
- P. praelongus Wulf. Shelburne Pond, (Pringle); Burlington, (N. F. F.). July.
- P. pusillus L. Malletts Bay, (Griffin). Aug.
- P. pusillus L., var. Sturrockii Benn. Malletts Bay, (Griffin). Aug.
- Y. Richardsonii (Benn.) Rydb. Quiet waters; occasional. July, Aug. Malletts Bay, (Griffin); Burlington, (N. F. F.).
- P. Robbinsii Oakes. Shelburne Pond, (Jones). July, Aug.
- P. zosterifolius Schum. Shelburne Pond, (Grout and Tracy, Jones); Malletts Bay, (Griffin). July, Aug.

JUNCAGINACEAE. ARROW GRASS FAMILY

SCHEUCHZERIA L.

S. palustris L. Peat bogs; rare. June. Colchester, (Torrey, Jones).

ALISMACEAE. WATER PLANTAIN FAMILY

ALISMA L. WATER PLANTAIN

A. Plantago-aquatica L. Swales; common. July, Aug.

SAGITTARIA L. ARROW-HEAD

- S. graminea Michx. Muddy shores of Lake Champlain and Winooski River; rare. July, Aug. Colchester, (Perkins, Griffin); Burlington, (N. F. F.).
- S. heterophylla Pursh. Low muddy shores; occasional. July, Aug.
- S. latifolia Willd. Wet places; common and presenting many different forms. Aug., Sept.

HYDROCHARITACEAE. FROG'S BIT FAMILY

ELODEA Michx. WATER-WEED

E. canadensis Michx. Slow streams and ponds; frequent. Aug.

VALLISNERIA (Mich.) L. TAPE GRASS. EEL GRASS

V. spiralis L. Slow water; frequent. Aug., Sept.

GRAMINEAE. GRASS FAMILY

AGROPYRON Gaertn.

- A. caninum (L.) Beauv. Awned Wheat Grass. Ledges at Winooski Gorge, Colchester, (Jones). July, Aug.
- A. REPENS (L.) Beauv. Couch, Quitch, or Quick Grass. Fields; common. June-Aug.
- A. tenerum Vasey. Slender Wheat Grass. Dry warm rocky woodlands and ledges; frequent. July, Aug.

AGROSTIS L. BENT GRASS

- A. alba L. Fiorin or White Bent Grass. Moist places; common and extremely variable. July. Var. vulgaris (With.) Thurb. Red Top. Fields and roadsides; common. July. Variable and intergrading with the species. A stoloniferous form (A. stolonifera auth. not L.) occurs in lawns, Burlington, (Jones). A small form occurs in cool pastures, sometimes having flowering glume awned; Burlington, (Jones).
- A. hyemalis (Walt.) BSP. Hair Grass. Dry ditches and other exsicated places; common. June-Aug.
- A. perennans (Walt.) Tuck. Thin Grass. Moist shaded places; frequent. Aug., Sept.

ALOPECURUS L. FOXTAIL GRASS

- A. geniculatus L., var. aristulatus Torr. Floating Foxtail Grass. Wet meadows; occasional. June, July. South Burlington, (Hinsdale); Burlington and Colchester, (Jones).
- A. PRATENSIS L. Meadow Foxtail Grass. Introduced in fields. May, June. Burlington, (Jones).

ANDROPOGON (Royen) L. BEARD GRASS

- A. furcatus Muhl. Forked Beard Grass. Dry rocky river banks; occasional. Aug., Sept.
- A. scoparius Michx. Broom Beard Grass. Dry sterile soil; occasional. Aug., Sept.

ANTHOXANTHUM L. SWEET VERNAL GRASS

A. ODORATUM L. Old meadows and roadsides; occasional. May-July.

ARISTIDA L. TRIPLE-AWNED GRASS

A. dichotoma Michx. Poverty Grass. Sterile gravelly soil; rare. Aug., Sept. Essex, (N. F. F.).

ARRHENATHERUM Beauv. OAT GRASS

A. ELATIUS (L.) Beauv. Tall Oat Grass. Meadows; rare. June, July. Burlington, (Jones).

BRACHYELYTRUM Beauv.

B. erectum (Schreb.) Beauv. Moist shaded places; frequent. July, Aug.

BROMUS L. BROME GRASS

- B. ciliatus L. Wood Chess. Moist thickets; common. July, Aug.
- B. INERMIS Leyss. Hungarian Brome Grass. Adventive in meadows, Experiment Farm, Burlington, (Orton); Colchester, (Griffin). July.
- B. Kalmii Gray. Wild Chess. Dry woodlands; occasional. June, July.
- B. purgans L. Thickets; frequent. July, Aug.
- B. SECALINUS L. Cheat or Chess. Fields and waste places; occasional. June-Aug.

CALAMAGROSTIS Adans. REED BENT GRASS.

C. canadensis (Michx.) Beauv. Blue-joint Grass. Marshes and wet places; common. June, July.

CENCHRUS L. SANDBUR. BUR GRASS

C. CAROLINIANUS Walt. Dry sandy soil; local. Aug., Sept. Colchester and Burlington, (N. F. F.); Essex, (Griffin). Evidently a recent introduction.

CINNA L. WOOD REED GRASS

- C. arundinacea L. Wood Reed Grass. Moist woods; rare. Aug., Sept. South Burlington, (Jones).
- C. latifolia (Trev.) Griseb. Slender Wood Reed Grass. Moist woods; rare. Aug., Sept. Burlington, (N. F. F.).

DACTYLIS L. ORCHARD GRASS

D. GLOMERATA L. Fields, generally in partial shade; common. June,
July. A proliferous form occurs, Burlington, (Jones, Morse).

DANTHONIA DC. WILD OAT GRASS

- D. compressa Aust. Dry woods; rare. June, July. South Burlington, (Jones).
- D. spicata (L.) Beauv. Common Wild Oat Grass. Dry sterile soil; common. June, July.

DESCHAMPSIA Beauv.

D. flexuosa (L.) Trin. Common Hair Grass. Dry places; common-June, July.

DIGITARIA Scop. FINGER GRASS

- D. HUMIFUSA Pers. Small Crab Grass. Occasional in sandy fields, more frequent as a lawn weed. Aug., Sept.
- D. SANGUINALIS (L.) Scop. Large Crab Grass. Cultivated fields and waste places; common. Aug., Sept.

ECHINOCHLOA Beauv.

E. CRUSGALLI (L.) Beauv. Barnyard Grass. Moist ground, waste places, etc.; common. Aug.-Oct. Presenting many differing forms; viz.: long awned, short awned, semi-prostrate, erect, etc.

ELYMUS L. WILD RYE. LYME GRASS

- F. canadensis L. Nodding Wild Rye. Low thickets and river banks; common. July, Aug. Var. GLAUCIFOLIUS (Muhl.) Gray. Dry banks; occasional. July, Aug.
- E. robustus Scribn. & Sm. Burlington, (Jones). July, Aug.
- E. striatus Willd. Slender Wild Rye. Ledges of Winooski River, (Pringle). July, Aug.
- E. virginicus L. Wild Rye. Moist thickets of river banks; frequent. July, Aug.

ERAGROSTIS Beauv.

- E. hypnoides (Lam.) BSP. Creeping Eragrostis. Gravelly or sandy shores; frequent. Aug., Sept.
- E, MEGASTACHYA (Koeler) Link. Strong-scented Eragrostis. Waste places; becoming frequent. July-Sept.
- E. PILOSA (L.) Beauv. Sandy soil, especially along railroads; rare. July-Sept. Burlington, (Priest, N. F. F.).

FESTUCA L. FESCUE GRASS

- F. ELATIOR L. Taller or Meadow Fescue. Meadows and roadsides; common. June, July.
- F. nutans Spreng. Nodding Fescue. Rocky woods; frequent. June, July.
- F. OVINA L. Sheep's Fescue. Introduced in lawn, South Burlington, (Jones). June, July.
- F. rubra L. Red Fescue. Rocky or sandy shores of Lake Champlain; occasional. June, July. Var. Subvillosa Mert. & Koch. Adventive in lawn, South Burlington, (Jones). July.

GLYCERIA R. Br. MANNA GRASS

- G. borealis (Nash) Batch. Floating Manna Grass. Wet places or in shallow water; frequent. June-Aug.
- G. canadensis (Michx.) Trin. Rattlesnake Grass. Wet places; common. July.
- G. grandis Wats. Reed Meadow Grass. Wet soil; common. July.
- G. nervata (Willd.) Trin. Manna Grass. Wet meadows; common. June, July.
- G. pallida (Torr.) Trin. Pale Manna Grass. Shallow water or wet soil; occasional. June-Aug.
- G. Torreyana (Spreng.) Hitchc. Wet woods; occasional. July, Aug.

HIEROCHLOE (Gmel.) R. Br. HOLY GRASS

H. odorata (L.) Wahlenb. Vanilla or Seneca Grass. Moist meadows; occasional. May, June.

HORDEUM (Tourn.) L. BARLEY

- H. AEGICERAS Nees. Egyptian Barley. Adventive in railroad yards, Burlington, (Jones, N. F. F.). July, Aug.
- H. JUBATUM L. Squirrel-tail Grass. Waste ground; occasional. June-Aug.

HYSTRIX Moench. BOTTLE-BRUSH GRASS

H. patula Moench. Moist woods; occasional. June, July.

LEERSIA Sw. CUT-GRASS. WHITE GRASS

- L. oryzoides (L.) Sw. Rice Cut-grass. Wet places; frequent. Aug., Sept.
- L. virginica Willd. White Grass. Wet woods; occasional. Aug.

LOLIUM L. DARNEL

- L. PERENNE L. Common Darnel, Perennial Ray or Rye Grass. Adventive by roadsides and in meadows; occasional. June, July.
- L. TEMULENTUM L. Bearded Darnel. Adventive in fields: rare. June, July. Essex, (Wild); Burlington, (Jones).

MELICA L. MELIC GRASS

M. striata (Michx.) Hitchc. Purple Oat. Rocky woods; frequent. May, June.

MUHLENBERGIA Schreb.

- M. mexicana (L.) Trin. Meadow Muhlenbergia. Moist meadows and banks; common. Aug., Sept. Variable.
- M. racemosa (Michx.) BSP. Marsh Muhlenbergia. Wet rocks and marshy places; frequent and variable. Aug., Sept.
- M. Schreberi Gmel. Drop-seed, Nimble Will. In shaded yard, Burlington, (Jones). Aug., Sept.
- M. tenuitlora (Willd.) BSP. Slender Muhlenbergia. Rocky woods; rare. Aug., Sept. Colchester, (Jones).

ORYZOPSIS Michx. MOUNTAIN RICE

- (), asperifolia Michx. White-grained Mountain Rice. Woods; common. June.
- O. pungens (Torr.) Hitchc. Slender Mountain Rice. Dry sandy soil; frequent. Apr., May.
- O. racemosa (Sm.) Ricker. Black-fruited Mountain Rice. Rocky woods; frequent. July, Aug.

PANICUM L. PANIC GRASS

- P. barbulatum Michx. Rocky woods and hillsides; rare. July. Colchester, (Jones).
- P. capillare L. Old-witch Grass. A common weed in sandy fields. Aug., Sept.
- P. clandestinum L. Moist thickets; frequent. June, July.
- P, depauperatum Muhl. Dry woods and fields; occasional. June-Aug.
- P. dichotomum L. Woods; rare. June, July. Burlington, (Jones).
- P. huachucae Ashe. Open ground; frequent. June, July.
- P. huachucae Ashe, var. silvicola Hitchc. & Chase. Woods; occasional. June, July.
- P. implicatum Scribn. Wet meadows; frequent. June, July.
- P. latifolium L. Woodlands; frequent. June, July.
- P. MILIACEUM L. European Millet. Waste places; rare. July, Aug. Burlington, (Brainerd, Jones).
- P. subvillosum Ashe. Dry woods and sandy ground. June, July. Burlington, (Jones).

- P. tennesseense Ashe. Moist ground; rare. June, July. Burlington. (N. F. F.).
- P. tsugetorum Nash. Sandy soil; frequent. June, July.
- P. Werneri Scribn. Sterile woods and knolls; rare. June-Aug. Burlington and Colchester, (Jones).
- P. xanthophysum Gray. Dry sandy soil; frequent. July, Aug. A broad-leaved form occurs, Burlington, (Jones).

PHALARIS L. CANARY GRASS

- P. arundinacea L. Reed Canary Grass. Wet grounds; occasional.

 June, July. Var. Picta L. Ribbon Grass. Persistent in old gardens and an occasional escape to roadsides. July.
- P. CANARIENSIS L. Canary Grass. Waste places; occasional, but probably only as seed is scattered from bird cages. June-Sept.

PHLEUM L.

P. PRATENSE L. Timothy, Herd's Grass. Meadows; common. June. July. Proliferous forms and plants bearing bulblets in the upper leaf axils occur, Burlington, (Jones). (See Vt. Exp. Sta. Bull. 95.)

PHRAGMITES Trin. REED

P. communis Trin. Wild Broom Corn. Swampy margins of ponds and lakes; occasional. Aug., Sept.

POA L. MEADOW GRASS. SPEAR GRASS

- P. alsodes Gray. Grove Spear Grass. Moist woods; frequent. May. June.
- P. ANNUA L. Low Spear Grass. Waysides and fields; common. Apr.-Sept.
- P. COMPRESSA L. Canada Blue Grass, Wire Grass. Dry fields and woodlands; common. May-Sept.
- P. debilis Torr. Weak Spear Grass. Dry rocky woodlands; frequent. May, June.
- P. nemoralis L. Rocky woodlands; occasional. June-Sept.
- P. prateusis L. June Grass, Spear Grass, Kentucky Blue Grass. Fields and meadows; common. May, June.
- P. triflora Gilib. Fowl Meadow Grass. Wet soil; common. July, Aug. The form known in Europe as P. FERTILIS Reich. occurs, Burlington, (Jones), probably introduced.

SETARIA Beauv. BRISTLY FOXTAIL GRASS

- S. GLAUCA (L.) Beauv. Yellow Foxtail, Pigeon Grass. Cultivated grounds and waste places; common. July-Sept.
- S. ITALICA (L.) Beauv. Millet, German Millet, Hungarian Grass. Occasionally spontaneous in cultivated ground. Aug., Sept.
- S. VERTICILLATA (L.) Beauv. Rough Foxtail Grass. Persistent in old garden, Burlington, (Jones). July, Aug.
- S. VIRIDIS (L.) Beauv. Green Foxtail, Bottle Grass. Cultivated grounds and waste places; common. July-Sept.

SORGHASTRUM Nash.

S. nutans (L.) Nash. Indian Grass, Wood Grass. Dry sandy soil; occasional. Aug., Sept.

SPARTINA Schreb. CORD OF MARSH GRASS

S. Michauxiana Hitche. Slough Grass. Shores of Lake Champlain and its tributaries; frequent. Aug., Sept.

SPHENOPHOLIS Scribn.

- s. nitida (Spreng.) Scribn. Woods; rare. June. Colchester, (Torrey).
- S. pallens (Spreng.) Scribn. Marshes; occasional. June, July.

SPOROBOLUS R. Br. DROP-SEED. RUSH GRASS

S. vaginiflorus (Torr.) Wood. Dry soil; occasional and variable. Sept.

TRISETUM Pers.

- T. melicoides (Michx.) Vasey. One form occurs. Var. MAJUS (Gray) Hitchc. Rocky banks of Winooski River, Colchester and Williston, (Pringle); Essex, (Jones). July.
- T. spicatum (L.) Richter. Narrow False Oat. Rocky banks; occasional. June,

ZIZANIA (Gronov.) L. WATER OF INDIAN RICE

- Z. aquatica L. Wild Rice, Water Oats. In shallow water; occasional. Aug., Sept.
- HORDEUM SATIVUM Jessen, Barley, Secale Cereale L., Rye, Triticum sativum Lam., Wheat and Zea Mays L., Indian Corn are occasional in waste places,

CYPERACEAE. SEDGE FAMILY

CAREX (Rupp.) L. SEDGE

- C. albicans Willd. Dry shady ledges; common. June, July.
- C. alopecoidea Tuckerm. Wet meadows; rare. June, July. Burlington, (Hazen).
- C. arctata Boott. Moist woodlands; common. May, June.
- C. aurea Nutt. Moist meadows and springy banks; frequent. June, July.
- C. Bebbii Olney. Low ground; rare. June, July. Burlington, (Hazen).
- C. bromoides Schkuhr. Bogs and swamps; frequent. June, July.
- C. brunnescens Poir. Open woods and dry rocky banks; rare. June, July. Colchester, (Griffin).
- C. canescens L., var. disjuncta Fern. Wet places; occasional. June, July.
- C. castanea Wahlenb. Low meadows; rare. June, July. Williston. (N. F. F.).
- C. cephaloidea Dewey. Moist woodlands; rare. June, July. South Burlington, (Hazen).
- C. cephalophora Muhl. Dry woods; frequent. June, July.
- C. chordorrhiza L. f. Cold bogs; rare. June, July. Colchester, (Jones, Grout).
- C. communis Bailey. Dry open woods; common. May, June.
- C. comosa Boott. Swamps; occasional. June, July.
- C. conoidea Schkuhr. Moist grassy places; occasional. June, July.
- C. Crawfordii Fern. Open sterile soil; frequent. June-Aug. Var. vicens Fern. Burlington, (Hazen). July.
- C. crinita Lam. Swales and along brooks; common. June, July.
- C. erinita Lam., var. gynandra (Schwein.) Schwein. & Torr. Same situations as the type; frequent. June, July.
- C. crinita Lam., var. minor Boott. Rare. June, July. Burlington, (Hazen).
- C. cristata Schwein. Wet grounds; occasional. June, July.

- C. debilis Michx., var. Rudgei Bailey. Moist sterile soil; rare. June, July. Colchester, (Hazen).
- C. Deweyana Schwein. Dry woodlands; common. June, July.
- C. diandra Schrank. Bogs; rare. June, July. South Burlington, (Hazen).
- C. digitalis Willd. Dry open woods; rare. June, July. Burlington, (Hazen).
- C. eburnea Boott. Dry shaded limestone; common. May-July.
- C. festucacea Schkuhr. Dry or rocky soil; rare. June, July. Burlington, (N. F. F.).
- C. filiformis L. Bogs and shallow water; occasional. June, July.
- C. flava L. Damp places; rare. June, July. Williston, (N. F. F.).
- C. foenea Willd. Dry woods and banks; rare. July. Colchester, (Griffin).
- C. folliculata L. Wet woods; rare. June-Aug. Burlington, (N. F. F.).
- C. formosa Dewey. Moist meadows; rare. June. Shelburne, (N. F. F).
- C. gracillima Schwein. Low meadows; common. June, July.
- C. granularis Muhl. Moist meadows; common. June, July.
- C. granularis Muhl., var. Haleana (Olney) Porter. Burlington, (Hazen). June, July.
- ('. grisea Wahlenb. Low meadows; rare. June. South Burlington, (N. F. F.).
- C. Houghtonii Torr. Dry sandy banks; rare. June, July. Burlington, (N. F. F.).
- C. hystericina Muhl. Swales; common. June, July.
- C. intumescens Rudge. Swamps and wet woods; occasional. June-Aug.
- ('. intumescens Rudge, var. Fernaldii Bailey. Same situations as the type; occasional. June-Aug.
- C. laxiflora Lam. Rich woods and meadows; occasional. May, June.
- ('. laxiflora Lam., var. patulifolia (Dewey) Carey. Burlington, (Jones).
 May, June.
- C. lenticularis Michx. Gravelly or sandy shores; rare. June-Aug. Shelburne, (Pringle).
- C. leptalea Wahlenb. Bogs and wet meadows; occasional. June, July.

- C. limosa L. Bogs; rare. June, July. Colchester and South Burlington, (Hazen).
- C. longirostris Torr. Rocky woods; rare. May, June. South Burlington, (Jones and Hazen); Colchester, (Griffin).
- C. lupuliformis Sartw. Marshes along Lake Champlain; occasional. July, Aug.
- C. lupulina Muhl. Swamps and ditches; common. July, Aug.
- C. lupulina Muhl., var. pedunculata Dewey. Swamps; rare. July, Aug. Burlington, (Hazen).
- C. lurida Wahlenb. Wet meadows; common. July, Aug. Very variable.
- C. mirabilis Dewey. Moist shady places; occasional. June, July.
- C. Muhlenbergii Schkuhr. Dry sandy soil; rare. June, July. Colchester, (Brainerd and Hazen).
- C. Oederi Retz., var. pumila (Coss. & Germ.) Fern. Shore of Lake Champlain, Shelburne, (Ross). June, July.
- C. pallescens L. Low meadows; common. June, July.
- C. pauciflora Lightf. Cold bogs; rare. June, July. Colchester, (Hazen).
- C. paupercula Michx., var. irrigua (Wahlenb.) Fern. Cold bogs; occasional. June-Aug.
- C. pedunculata Muhl. Rich woods; frequent. May.
- C, pennsylvanica Lam. Dry sandy soil; common. May, June.
- C. pennsylvanica Lam., var. lucorum (Willd.) Fern. Richer damper soil than the type. May, June. Burlington, (Brainerd).
- C. plantaginea Lam. Rich woods; frequent. May.
- C. platyphylla Carey. Shady banks; common. May, June.
- C. prasina Wahlenb. Springy bogs in woods; occasional. June, July.
- C. pubescens Muhl. Copses and rich woods; rare. May, June. South Burlington, (Shaw).
- C. retrorsa Schwein. Swamps; common. June, July.
- C. riparia Curtis. Swales; occasional. June, July.
- C. rosea Schkuhr. Dry open woods; frequent. June, July.
- C. rostrata Stokes, var. utriculata (Boott) Bailey. Wet swamps; occasional. June, July.

- C. scabrata Schwein. Springy bogs in woods; frequent. June, July.
- C. scirpoides Schkuhr. Boggy meadows; frequent. June, July.
- C. scoparia Schkuhr. Low grounds; common. June, July. Var. con-DENSA Fern. Colchester, (Hazen). June, July. Var. MONILIFORMIS Tuck. Burlington, (Hazen). June, July.
- C. siccata Dewey. Sandy shore of Lake Champlain; rare. June, July. Burlington, (Hazen).
- C. sparganioides Muhl. Rich woods; occasional. June, July.
- C. stellulata Good. Low open ground; frequent. June, July.
- C. stellulata Good., var. angustata Carey. Wet ground; rare. June, July. South Burlington, (N. F. F.).
- C. stellulata Good., var. excelsior (Bailey) Fern. Low open ground; frequent. June, July.
- C. stipata Muhl. Swales; common. June, July.
- C. straminea Willd. Dryish copses and fields; frequent. June, July.
- C. straminea Willd., var. echinodes Fern. Moist sandy soil, Burlington, (Brainerd). July.
- C. stricta Lam. Swales and marshes; frequent. June, July.
- C. tenella Schkuhr. Cold swamps; occasional. May, June.
- C. tenuiflora Wahlenb. Bogs; rare. June, July. Burlington, (Robbins).
- C. torta Boott. By streams; rare. June. Colchester, (Griffin).
- C. tribuloides Wahlenb. Open swales; frequent. July, Aug.
- C. tribuloides Wahlenb., var. reducta Bailey. Moist copses; frequent. July, Aug.
- C. trisperma Dewey. Cold bogs in shade; occasional. June, July.
- C. Tuckermani Dewey. Swales and marshes; frequent. June, July.
- C. typhinoides Schwein. (C. typhina Michx.) Low margin of Lake Champlain, Burlington, (N. F. F.). July, Aug.
- C. umbellata Schkuhr. Dry sandy soil; occasional. May, June.
- C. varia Muhl. Dry woods; rare. May, June. Colchester, (Brainerd and Jones); Burlington, (Pringle).
- C. vesicaria L. Two forms occur. Var. Jejuna Fern. Swales; rare. June, July. Burlington, (Hazen). Var. MONILE (Tuck.) Fern. Swales; common. June, July.

C. vulpinoidea Michx. Moist pastures; common. June, July.

CYPERUS (Tourn.) L. GALINGALE

- C. aristatus Rottb. Awned Cyperus. Wet sandy soil; frequent. Aug., Sept.
- C. dentatus Torr. Sandy shores of Lake Champlain; rare. Aug., Sept. Shelburne, (Pringle); Burlington and Colchester, (N. F. F.). Spikelets often abortive and changed into leafy tufts.
- C. esculentus L. Yellow Nut Grass. Lake shore and river banks; occasional. Aug., Sept.
- C. filiculmis Vahl, var. macilentus Fern. Dry sandy soil; common. July, Aug.
- C. Houghtonii Torr. On dry shifting sand, Colchester, (N. F. F.). Aug., Sept.
- C. rivularis Kunth. Shining Cyperus. Lake shore and river banks; frequent. Aug., Sept.
- C. strigosus L. Straw-colored Cyperus. Moist soil; frequent. July-Sept.

DULICHIUM Pers.

D. arundinaceum (L.) Britton. Borders of ponds; occasional. July-Sept.

ELEOCHARIS R. Br. SPIKE RUSH

- E. acicularis (L.) R. & S. Muddy shores below high water mark; common. July-Sept.
- E. diandra Wright. Wet sands of Winooski River; rare. Sept., Oct. Burlington, (Brainerd, N. F. F.).
- E. intermedia (Muhl.) Schultes. Bank of Winooski River, Burlington, (Brainerd, N. F. F.). Sept., Oct.
- E. obtusa (Willd.) Schultes. Muddy places; common. July-Sept.
- E. palustris (L.) R. & S. Wet ground; frequent. June-Aug. Var. GLAUCESCENS (Willd.) Gray. Moist grassy places; frequent. July, Aug. Var. vigens Bailey. (Var. major Sonder) Mouth of Winooski River, (Grout, Brainerd). July, Aug.
- E. tenuis (Willd.) Schultes. Wet meadows; rare. June-Aug. South Burlington, (N. F. F.).

ERIOPHORUM L. COTTON GRASS

- E. callitrix Cham. Hare's Tail. Peat bogs; frequent. May, June.
- E. tenellum Nutt. Swamps and bogs; rare. July, Aug. Colchester, (N. F. F.).
- E. virginicum L. Cotton Grass. Bogs and wet meadows; common. Aug., Sept.
- E. viridi-carinatum (Engelm.) Fern. Bogs; frequent. June, July.

RYNCHOSPORA Vahl. BEAK RUSH

- R. alba (L.) Vahl. White Beak Rush. Bogs; frequent. July-Sept.
- R. fusca (L.) Ait. f. Brown Beak Rush. Burlington, (Torrey). July-Sept. One specimen in U. V. M. Herb.

SCIRPUS (Tourn.) L. BULRUSH OF CLUB RUSH

- S. americanus Pers. Chair-maker's Rush. Sandy shores of Lake Champlain; frequent. July-Sept.
- S. atrocinctus Fern. Wet meadows; occasional. July. Var. BRACHY-PODUS Fern. Rare. July. Colchester, (N. F. F.).
- S. atrovirens Muhl. Moist meadows; common. July, Aug.
- S. cyperinus (L.) Kunth. Wool Grass. Wet meadows; common. Aug., Sept.
- S. fluviatilis (Torr.) Gray. River Bulrush. Marshy borders of bays and tributaries of Lake Champlain; occasional. July-Sept.
- S. georgianus Harper. Moist meadows; frequent. July, Aug. Proliferous forms occur, Burlington and Colchester, (N. F. F.).
- S. heterochaetus Chase. Sheltered shores; frequent. July, Aug.
- S. pedicellatus Fern. Swales; frequent. July, Aug.
- S. rubrotinctus Fern. Wet meadows; frequent. July, Aug.
- S. Smithii Gray. Shores of Lake Champlain. Aug. Abundant at one station, Burlington, (Grout and Tracy).
- S. validus Vahl. Great Bulrush. In shallow water; frequent. July, Aug.

ARACEAE. ARUM FAMILY

ACORUS L. SWEET FLAG. CALAMUS

A. Calamus L. Marshes; common. June.

ARISAEMA Martius. Indian Turnip. Dragon Arum

A. triphyllum (L.) Schott. Indian or Wild Turnip, Jack-in-the-Pulpit. Rich moist woods; common. Apr., May. Spathes, petioles and sheaths vary in color from pale green to dark purple with whitish stripes or spots.

CALLA L. WATER ARUM

- C. palustris L. Wild Calla. Bogs and marshes; frequent. May, June.

 SYMPLOCARPUS Salisb. SKUNK CABBAGE
- S. foetidus (L.) Nutt. Wet places; common. Apr., May.

LEMNACEAE. DUCKWEED FAMILY

LEMNA L. DUCKWEED. DUCK'S MEAT

- L. minor L. Lesser Duckweed. Stagnant water; frequent. Not found in flower.
- L. trisulca L. Ivy-leaved Duckweed. Ponds; occasional. Not found in flower,

SPIRODELA Schleid.

S. polyrhiza (L.) Schleid. Greater Duckweed. Stagnant water; frequent. Not found in flower.

ERIOCAULACEAE. PIPEWORT FAMILY

ERIOCAULON (Gronov.) L. PIPEWORT

E. articulatum (Huds.) Morong. (E. septangulare With.) Border of Lake Champlain, Colchester, (N. F. F., Griffin). Aug., Sept.

PONTEDERIACEAE. PICKEREL-WEED FAMILY

HETERANTHERA R. & P. MUD PLANTAIN

H. dubia (Jacq.) MacM. Water Star Grass. Ponds and streams; rare. Aug. Mouth of Winooski River, (Grout and Tracy). A small form occurs on muddy shore, Burlington, (N. F. F.).

PONTEDERIA L. PICKEREL-WEED

P. cordata L. Borders of ponds and slow streams; frequent. July, Aug.

JUNCACEAE. RUSH FAMILY

JUNCUS (Tourn.) L. RUSH. BOG RUSH

- J. acuminatus Michx. Sharp-fruited Rush. Wet meadows; rare. June-Aug. Essex, (Jones). This was a proliferous form of the species.
- J. alpinus Vill., var. insignis Fries. Wet gravelly shores of Lake Champlain; rare. July, Aug. Shelburne, (Pringle); Burlington, (N. F. F.).
- J. articulatus L. Jointed Rush. Wet grounds; rare. July, Aug. Essex, (Jones); Colchester, (N. F. F.).
- J. brevicaudatus (Engelm.) Fern. Moist places; frequent. July-Sept.
- J. bufonius L. Toad Rush. Low grounds; common. June-Sept.
- J. Dudleyi Wiegand. Damp grounds; occasional. June-Sept.
- J. effusus L. Bog Rush. Marshy grounds; common. July, Aug.
- J. nodosus L. Knotted Rush. Swamps and gravelly banks; frequent. July, Aug.
- J. pelocarpus Mey. Brownish-fruited Rush. Shores of Lake Champlain; rare. Aug., Sept. Shelburne, (Pringle); Colchester, (N. F. F.).
- J. tenuis Willd. Slender Rush. Fields; common and variable. June-Sept.

LUZULA DC. WOOD RUSH

- L. campestris (L.) DC., var. multiflora (Ehrh.) Celak. Common Wood Rush. Meadows and open woods; frequent. May, June.
- L. saltuensis Fern. Hairy Wood Rush. Woods and banks; frequent. May.

LILIACEAE. LILY FAMILY

ALLIUM (Tourn.) L. ONION. GARLIC

A. tricoccum Ait. Wild Leek. Rich woods; rare. June, July. Colchester, (Griffin).

ASPARAGUS (Tourn.) L. ASPARAGUS

A. officinalis L. Garden Asparagus. Roadside escape from gardens; frequent. June,

CLINTONIA Raf.

C. borealis (Ait.) Raf. Yellow Clintonia. Cool moist woods; common. May, June.

CONVALLARIA L. LILY OF THE VALLEY

C. Majalis L. Persistent near old building site, Shelburne, (N. F. F.). May, June.

ERYTHRONIUM L. DOG'S-TOOTH VIOLET

E. americanum Ker. Yellow Adder's Tongue. Rich copses and woodlands; common. Apr., May.

HEMEROCALLIS L. DAY LILY

H. FULVA L. Common Day Lily. Roadside escape from gardens; occasional. July.

LILIUM (Tourn.) L. LILY

- L. canadense L. Meadow Lily, Wild Yellow Lily. Moist meadows; common. July.
- L. philadelphicum L. Wood Lily, Wild Orange-red Lily. Sandy open woodlands: common. July.
- L. TIGRINUM Ker. Tiger Lily. Roadside escape from gardens; rare. Aug. Burlington, (Miss Brown); Essex, (N. F. F.).

MAIANTHEMUM Wiggers.

M. canadense Desf. Two-leaved Solomon's Seal, Wild Lily of the Valley. Woods; common. May, June.

MEDEOLA (Gronov.) L. INDIAN CUCUMBER-ROOT

M. virginiana L. Rich moist woods; common. May, June.

OAKESIA Wats.

 sessilifolia (L.) Wats. Sessile-leaved Bellwort. Moist woodlands; common. May, June.

POLYGONATUM (Tourn.) Hill. SOLOMON'S SEAL

P. biflorum (Walt.) Ell. Small Solomon's Seal. Wooded hillsides; common. May, June.

SMILACINA Desf. FALSE SOLOMON'S SEAL

- S. racemosa (L.) Desf. False Spikenard. Moist copses; common.

 May, June.
- S. stellata (L.) Desf. Star-flowered Solomon's Seal. Moist banks and meadows; frequent. May.
- S. trifolia (L.) Desf. Three-leaved Solomon's Seal. Cold sphagnum swamps; occasional. June, July.

SMILAX (Tourn.) L. CAT BRIER. GREEN BRIER

S. herbacea L. Carrion-flower. River banks and moist thickets; occasional. June.

STREPTOPUS Michx. TWISTED-STALK

S. roseus Michx. Cold damp woods; frequent. May.

TRILLIUM L. WAKE ROBIN, BIRTHROOT

- T. cernuum L. Nodding Trillium. Moist woods; rare. May. South Burlington, (Hood).
- T. erectum L. Purple Trillium. Rich woods; frequent. May.
- T. grandiflorum (Michx.) Salisb. White Trillium. Rich moist woods; common. May.
- T. undulatum Willd. Painted Trillium. Cool moist woods; occasional.

 May.

UVULARIA L. BELLWORT

U. grandiflora Sm. Large-flowered Bellwort. Rich woods; frequent. Apr., May. A form occurs the leaves of which are variegated with white; Shelburne, (Mrs. Gebhardt).

VERATRUM (Tourn.) L. FALSE HELLEBORE

V. viride Ait. American White Hellebore, Indian Poke. Wet meadows and swamps; common. June.

IRIDACEAE. IRIS FAMILY

IRIS (Tourn.) L. FLEUR-DE-LIS

- I. PSEUDACORUS L. Yellow Iris. A garden escape, bank of Winooski River, Essex; (Ross). June.
- J. versicolor L. Wild Iris, Blue Flag. Wet meadows; common. June, July.

SISYRINCHIUM L. BLUE-EYED GRASS

S. angustifolium Mill. Meadows, fields and damp sandy soil; common. May, June.

ORCHIDACEAE. ORCHIS FAMILY

ARETHUSA (Gronov.) L.

A. bulbosa L. Sphagnum bogs; occasional. June. A white-flowered form occurs, South Burlington, (Howe).

CALOPOGON R. Br.

C. pulchellus (Sw.) R. Br. Grass Pink. Bogs; occasional. June, July.

CORALLORRHIZA (Haller) Chat. CORAL ROOT

- C. maculata Raf. Large Coral Root. Woods; occasional. July, Aug.
- C. trifida Chat. Early Coral Root. Swamps and damp woods; occasional. May, June.

CYPRIPEDIUM L. LADY'S SLIPPER. MOCCASIN FLOWER

- C. acaule Ait. Stemless or Pink Lady's Slipper. Dry woods; frequent.

 May. June.
- C. arietinum R. Br. Ram's Head Lady's Slipper. Occasional in rocky woods where it reaches great perfection, colonies of thirty-nine to fifty blossoms having been seen. May, June.
- C. hirsutum Mill. Showy Lady's Slipper. Cold swamps; occasional. June, July.
- C. parviflorum Salisb. Smaller Yellow Lady's Slipper. Bogs and low woods; rare. May, June. South Burlington, (Torrey); Colchester, (Griffin).
- C. parviflorum Salisb., var. pubescens (Willd.) Knight. Large Yellow Lady's Slipper. Rich moist woods; frequent. May, June.

EPIPACTIS (Haller) Boehm. RATTLESNAKE PLANTAIN

- E. pubescens (Willd.) Eaton. Downy Rattlesnake Plantain. Dry woods; occasional. Aug.
- E. repens (L.) Crantz, var. ophioides (Fern.) Eaton. Lesser Rattle-snake Plantain. Cold mossy woods; rare. July. South Burlington, (Mrs. Walker).
- E. tesselata (Lodd.) Eaton. Cold moist woods; rare. Aug. Burlington, (Collins); South Burlington, (N. F. F.).

HABENARIA Willd. REIN ORCHIS. FRINGED ORCHIS

- H. blephariglottis (Willd.) Torr. White Fringed Orchis. Bogs; rare.
 Aug. South Burlington, (Torrey).
- H. bracteata (Willd.) R. Br. Long-bracted Orchis. Damp woods and meadows; occasional. May, June.
- H. clavellata (Michx.) Spreng. Small Wood Orchis. Bogs; rare. July, Aug. South Burlington, (Torrey, Jones); Burlington, (Ross, N. F. F.).
- H. dilatata (Pursh) Gray. White Bog Orchis. Cold bogs; rare. June,July. South Burlington, (Hazen).
- H. fimbriata (Ait.) R. Br. Large Purple Fringed Orchis. Cool moist woods; rare. June, July. Essex, (Jones and Hazen).
- H. Hookeri Torr. Dry woods; occasional. June.
- H. hyperborea (L.) R. Br. Tall Green Orchis. Bogs and cold woods; occasional. July.
- H. lacera (Michx.) R. Br. Ragged Fringed Orchis. Moist ground; rare. July, Aug. South Burlington, (Torrey); Burlington, (N. F. F.).
- H. macrophylla Goldie. Large-leaved Orchis. Moist coniferous woods; rare. June, July. Williston, (Bates); South Burlington, (Ross).
- II. orbiculata (Pursh) Torr. Round-leaved Orchis. Cold rich woods; rare, June, July. Burlington, (N. F. F., H. A. P. Torrey).
- H. psycodes (L.) Sw. Smaller Purple Fringed Orchis. Wet meadows and bogs; occasional. July, Aug.

LIPARIS Richard. TWAYBLADE

L. liliifolia (L.) Richard. Large Twayblade. Woods; rare. June, July. Colchester, (Eggleston). Specimen lost.

I. Loeselii (L.) Richard. Fen Orchis. Damp fields; occasional. June, July.

LISTERA R. Br. TWAYBLADE

L. cordata (L.) R. Br. Heart-leaved Twayblade. Sphagnum bogs; rare. June. South Burlington, (Jones).

MICROSTYLIS (Nutt.) Eaton. ADDER'S MOUTH

M. unifolia (Michx.) BSP. Green Adder's Mouth. Open woods; occasional. July.

ORCHIS (Tourn.) L.

 spectabilis L. Showy Orchis. Rich moist woods; occasional. May, June.

POGONIA Juss.

- P. affinis Aust. Small Whorled Pogonia. One plant in woods at Fairholt, Burlington, (Mrs. Holt). May, June.
- P. ophioglossoides (L.) Ker. Rose Pogonia, Snake-mouth. Bogs; frequent. June, July.
- P. verticillata (Willd.) Nutt. Whorled Pogonia. Woods; rare. May, June. Colchester, (Robbins, Torrey, Ross); Burlington, (Jones). Has been found in two or three different stations at intervals of several years.

SPIRANTHES Richard. LADIES' TRESSES

- S. cernua (L.) Richard. Nodding Ladies' Tresses. Wet soil; commou. Aug., Sept.
- S. gracilis (Bigel.) Beck. Slender Ladies' Tresses. Sandy woods and plains; occasional. July, Aug.
- S. Romanzoffiana Cham. Swamps and moist soil; rare. July. Aug. Colchester, (Griffin).

SALICACEAE. WILLOW FAMILY

POPULUS (Tourn.) L. POPLAR. ASPEN

- P. ALBA L. White Poplar, Silver-leaved Poplar, Abele. Frequent as an escape from cultivation. Apr., May. Only pistillate trees occur.
- P. balsamifera L. Balsam Poplar, Tacamahac. Wet ravines and borders of rivers; frequent. Apr., May.

- P. CANDICANS Ait. Balm of Gilead. Occasional in cultivation and sometimes spontaneous. Apr., May.
- P. deltoides Marsh. Cotton-wood, Necklace Poplar. Borders of streams and of Lake Champlain; frequent. Apr., May.
- P. grandidentata Michx. Large-toothed Aspen. Rich woods; frequent. Apr., May.
- P. NIGRA L. One form occurs. Var. ITALICA Du Roi. Lombardy Poplar. Frequently cultivated and tends to spread by sprouting from the roots. Apr., May. Only staminate trees occur.
- P. tremuloides Michx. American Aspen, Trembling Poplar. Woods; common. Apr., May.

SALIX (Tourn.) L. WILLOW. OSIER

- S. ALBA L. One form occurs. Var. VITELLINA (L.) Koch. White Willow. Common in cultivation and occasional as an escape. Apr., May.
- S. cordata Muhl. Heart-leaved Willow. Wet soil; common. Apr., May.
- S. discolor Muhl. Pussy or Glaucous Willow. Wet soil; common. Apr., May. Var. eriocephala (Michx.) Anders. South Burlington, (Kirk and Jones). Apr., May.
- S. FRAGILIS L. Crack Willow. Common in cultivation and as an escape. May,
- S. humilis Marsh. Prairie Willow. Dry sandy plains; common. Apr., May.
- S. longifolia Muhl. Sand-bar Willow. Shores of Lake Champlain; occasional. May.
- S. lucida Muhl. Shining Willow. Moist banks; frequent. May.
- S. nigra Marsh. Black Willow. Banks of streams and ponds; common. May. Var. FALCATA (Pursh) Torr. Burlington, (Rand). May.
- S. pedicellaris Pursh. Bog Willow. Peat bogs; frequent. May.
- S. petiolaris Sm. Slender Willow. Swamps and shores of Lake Champlain; rare. May. Burlington, (Grout and Jones).
- S. PURPUREA L. Purple Willow. A frequent escape to watersides. Apr., May.

- S. rostrata Richards. Beaked Willow. Borders of thickets; common. Apr., May.
- **S. sericea Marsh.** Silky Willow. In swamps and along streams; rare. Apr., May. Burlington and South Burlington, (Jones).
- Some peculiar willows occur at Burlington, (Jones), which will probably prove to be hybrids,

MYRICACEAE. SWEET GALE FAMILY

MYRICA L.

- M. asplenifolia L. Sweet Fern. Dry sterile soil; common. Apr., May.
- M. Gale L. Sweet Gale. Swamps and borders of ponds; occasional. May.

JUGLANDACEAE. WALNUT FAMILY

CARYA Nutt. HICKORY

- C. alba (L.) Koch. Mocker Nut, White-heart Hickory. Rocky pasture, Burlington, (Burns). June.
- C. cordiformis (Wang.) Koch. Bitter-nut or Swamp Hickory. Occasional in cultivation, but rare otherwise. May, June. South Burlington, (Hubbard and Strait).
- C. ovata (Mill.) Koch. Shellbark or Shagbark Hickory, Walnut. Rich soil; frequent. May, June.

JUGLANS L. WALNUT

J. cinerea L. Butternut, White Walnut. Rocky hillsides; frequent. May.

BETULACEAE. BIRCH FAMILY

ALNUS (Tourn.) Hill. ALDER

- A. incana (L.) Moench. Speckled or Hoary Alder. Swamps and borders of streams; common. Apr. Several forms occur, which are intermediate between this and A. rugosa, and are regarded by some as hybrids.
- A. mollis Fern. Downy Green Alder. Damp thickets and exposed rocky banks; rare. May. Burlington, (N. F. F.).

A. rugosa (Du Roi) Spreng. Smooth Alder. Moist soil; frequent. Apr.

BETULA (Tourn.) L. BIRCH

- B. alba L., var. papyrifera (Marsh.) Spach. Paper, Canoe or White Birch. Moist soil; common. Apr., May.
- B. lenta L. Cherry, Sweet or Black Birch. Rich woodlands; frequent. Apr., May.
- B. luten Michx. f. Yellow or Gray Birch. Rich cool soils; frequent. Apr., May.
- B. populifolia Marsh. White, Gray or Old Field Birch. Poor sandy soil; common. Apr., May.

CARPINUS (Tourn.) L. HORNBEAM. IRONWOOD

C. caroliniana Walt. American Hornbeam, Blue or Water Beech. Along streams; frequent. May.

CORYLUS (Tourn.) L. HAZELNUT, FILBERT

C. rostrata Ait. Beaked Hazelnut. Dry thickets; common. Apr., May.

OSTRYA (Mich.) Scop. HOP HORNBEAM. IRON WOOD

 virginiana (Mill.) Koch. American Hop Hornbeam, Leverwood. Rich woods; common. May.

FAGACEAE. BEECH FAMILY

CASTANEA (Tourn.) Hill. CHESTNUT

C. dentata (Marsh.) Borkh. Chestnut. Rocky woods and hillsides.

July. The trees on the Ethan Allen farm in the north part of

Burlington were supposed to represent the northern limit of this

tree, but it has recently been found in Colchester, (Griffin).*

FAGUS (Tourn.) L. BEECH

F. grandifolia Ehrh. American Beech. Rich uplands; common. May.

QUERCUS (Tourn.) L. OAK

- Q. alba L. White Oak. Dryish woods; common. May.
- Q. bicolor Willd. Swamp White Oak. Borders of streams and swamps; frequent. May.

^{*}This last station is probably the one referred to in Hemenway's Vermont Gazeteer, Vol. 1, p. 758.

- Q. macrocarpa Michx. Bur Oak, Over-cup or Mossy-cup Oak. Rich soil; rare, May. Burlington, (Jones); Colchester, (Griffin).
- Q. Muhlenbergii Engelm. Yellow Oak, Chestnut Oak. Dry limestone hillsides; rare. May. Colchester, (Griffin).
- Q. rubra L. Red Oak. Rich or poor soil; common. May. Var. AMBIGUA (Michx. f.) Fern. Gray Oak. Rocky woods, Burlington, (Burns). May.
- Q. velutina Lam. Quercitron, Yellow-barked or Black Oak. Dry light soil; frequent. May.

URTICACEAE. NETTLE FAMILY

BOEHMERIA Jacq. FALSE NETTLE

- B. cylindrica (L.) Sw. Moist shady ground; frequent. July, Aug.

 CANNABIS (Tourn.) L. HEMP
- C. SATIVA L. Common Hemp. Waste places; occasional. July-Sept.

 CELTIS (Tourn.) L. Nettle Tree. Hackberry
- C. occidentalis L. Sugarberry. River banks and intervals; occasional.

 May.

HUMULUS L. HOP

- H. JAPONICUS Sieb. & Zucc. Japanese Hop. Waste places; rare. July. Burlington, (Jones, N. F. F.).
- H. Lupulus L. Common Hop. An occasional roadside escape. July.

LAPORTEA Gaud. WOOD NETTLE

L. canadensis (L.) Gaud. Rich woods; frequent. July, Aug.

MORUS (Tourn.) L. MULBERRY

M. Alba L. White Mulberry. An occasional escape to woods and roadsides. May. Burlington, (N. F. F.).

PARIETARIA (Tourn.) L. PELLITORY

P. pennsylvanica Muhl. Shaded rocky banks; rare. June, July. Burlington, (Jones); South Burlington, (N. F. F.); Colchester, (Griffin).

PILEA Lindl. RICHWEED. CLEARWEED

P. pumila (L.) Gray. Cool and moist shaded places; frequent. July, Aug.

ULMUS (Tourn.) L. ELM

- U. americana L. American or White Elm. Moist soil; common. Apr.
- U. CAMPESTRIS L. English Elm. Commonly planted and escaped to roadside, Colchester, (N. F. F.). Apr.
- U. fulva Michx. Slippery or Red Elm. Rich rocky woods; occasional. Apr.
- U. racemosa Thomas. Cork or Rock Elm. Occurs as a shade tree in Burlington and search will probably reveal it on the river banks or calcareous ridges in this vicinity. Apr., May.

URTICA (Tourn.) L. NETTLE

- U. gracilis Ait. Slender or Tall Wild Nettle. Fence rows and moist ground; common. July, Aug.
- U. Lyallii Wats. Alluvial thickets and waste places; frequent. July, Aug. Has hitherto been confused with U. graeilis.

SANTALACEAE. SANDALWOOD FAMILY

COMANDRA Nutt. BASTARD TOAD-FLAX

- C. Richardsiana Fern. Dry sandy or gravelly soil; common. May, June.
- C. umbellata (L.) Nutt. Dry ground; rare. May, June. Colchester, (Griffin).

LORANTHACEAE. MISTLETOE FAMILY

ARCEUTHOBIUM Bieb.

A. pusillum Peck. Dwarf Mistletoe. Parasitic on spruce and tamarack. May. Colchester, (Jones).

ARISTOLOCHIACEAE. BIRTHWORT FAMILY

ASARUM (Tourn.) L. ASARABACCA. WILD GINGER

A. canadense L. Rich woods; common. Apr., May.

POLYGONACEAE. BUCKWHEAT FAMILY

FAGOPYRUM (Tourn.) Hill. BUCKWHEAT

F. ESCULENTUM Moench. Persisting after cultivation. June-Sept.

POLYGONELLA Michx.

P. articulata (L.) Meisn. Coast Jointweed. Sandy shore of Lake Champlain; rare. Aug., Sept. Colchester, (Pringle, N. F. F.).

POLYGONUM (Tourn.) L. KNOTWEED

- P. acre HBK., var. leptostachyum Meisn. Water Smartweed. Wet meadows; occasional. July-Sept.
- P. amphibium L. One form occurs. Var. Hartwrightii (Gray) Bissell. Wet places; rare. Aug., Sept. Burlington, (Jones, N. F. F.).
- P. arifolium L. Halberd-leaved Tear-thumb. Wet ground; occasional. Aug., Sept.
- P. aviculare L. Knotgrass, Knotweed, Doorweed. Dooryards, waste places, etc.; common. June-Sept.
- P. aviculare L., var. vegetum Ledeb. Burlington, (Grout). July-Sept.
- P. cilinode Michx. Fringed Black Bindweed. Rocky copses; frequent. July-Sept.
- P. Convolvulus L. Black Bindweed. Cultivated and waste ground; common. June-Sept:
- P. CUSPIDATUM Sieb. & Zucc. Japanese Knotweed. Adventive in garden, Shelburne, (Mrs. Gebhardt). Aug., Sept.
- P. erectum L. Erect Knotweed. Waysides, waste places, etc.; local. July, Aug. South Burlington, (N. F. F.).
- P. Hydropiper L. Common Smartweed or Water Pepper. Moist places; common. July-Sept.
- P. lapathifolium L. Dock-leaved or Pale Persicaria. Alluvial fields; frequent. July-Sept. A stout form near var. NODOSUM (Pers.) Weinm. occurs, Fort Ethan Allen, (N. F. F.).
- P. Muhlenbergii (Meisn.) Wats. Swamp Persicaria. Muddy or exsicated places; occasional. Aug., Sept.
- P. ORIENTALE L. Prince's Feather. Persisting about gardens; occasional. July-Sept.

- P. pennsylvanicum L. Common or Pink Persicaria. Moist soil; frequent. July-Sept.
- P. Persicaria L. Lady's Thumb. Waste places; common. July-Sept.
- P. sagittatum L. Arrow-leaved Tear-thumb. Low ground; common. Aug., Sept.
- P. scandens L. Climbing False Buckwheat. Moist thickets; frequent. Aug., Sept.
- P. virginianum L. Moist thickets; occasional. Aug., Sept.

RUMEX L. DOCK. SORREL

- R. ACETOSA L. Garden Sorrel. Adventive in grass lands, Burlington, (Jones). June.
- R. ACETOSELLA L. Field or Sheep Sorrel. Fields, especially in sterile soil; a common weed. May-July.
- R. Britannica L. Great Water Dock. Wet places; frequent. July, Aug.
- R. CRISPUS L. Yellow Dock. Fields and waste places; common. June, July.
- R. MEXICANUS Meisn. Rich soil; rare. July. Colchester, (Sprague); Burlington, (N. F. F.).
- R. MONTANUM Desf. A garden escape, Burlington, (N. F. F.). June. Station destroyed by building operations.
- R. OBTUSIFOLIUS L. Bitter Dock. Fields and waste places; common. June, July.
- R. PATIENTIA L. Patience Dock. Rich open soil; frequent. June, July.
- R. verticillatus L. Swamp Dock. Swamps; frequent. June, July.

CHENOPODIACEAE. GOOSEFOOT FAMILY

CHENOPODIUM (Tourn.) L. GOOSEFOOT. PIGWEED

- C. ALBUM L. Lamb's Quarters, Pigweed. Fields; a common weed. June-Sept. Var. viride (L.) Moq. Becoming frequent. June-Sept.
- C. Ambrosioides L. Mexican Tea. Adventive on railway embankment, Colchester, (N. F. F.). Aug., Sept.
- C. Botrys L. Jerusalem Oak, Feather Geranium. Lake shore and waste places; occasional. July-Sept.

- C. capitatum (L.) Asch. Strawberry Blite. Rich ground; rare. July. Burlington, (Jones and Howe).
- C. GLAUCUM L. Oak-leaved Goosefoot. Railroad yard, Burlington, (Orton). July-Sept.
- C. hybridum L. Maple-leaved Goosefoot. Shady ledges and waste places; frequent. July-Sept.
- C. URBICUM L. Upright Goosefoot. Adventive in lumber yards, Burlington, (Howe). June-Aug.

SALSOLA L. SALTWORT

S. KALI L., var. TENUIFOLIA Mey. Russian Thistle. Occasional as a recent introduction, Burlington, (N. F. F.); Colchester, (Jones). Aug., Sept.

AMARANTHACEAE. AMARANTH FAMILY

ACNIDA L. WATER HEMP

A. tuberculata Moq. Sandy shore of Lake Champlain, Burlington, (N. F. F.). Aug., Sept.

AMARANTHUS (Tourn.) L. AMARANTH

- A. BLITOIDES Wats. Prostrate Amaranth. Waste places and along railroads; becoming common. July-Sept.
- A. GRAECIZANS L. Tumble Weed. Waste and cultivated ground; frequent. July-Sept.
- A. RETROFLEXUS L. Green Amaranth, Pigweed. A common weed in cultivated ground. July-Sept.
- A. SPINOSUS L. Thorny Amaranth. Adventive in cultivated ground, Burlington, (Mrs. Walker). July-Sept.

PHYTOLACCACEAE. POKEWEED FAMILY

PHYTOLACCA (Tourn.) L. POKEWEED

P. decandra L. Common Poke or Scoke, Garget, Pigeon Berry. Pastures and fields; rare. July-Sept. Colchester, (Jones).

AIZOACEAE

MOLLUGO L. INDIAN CHICKWEED

M. VERTICILLATA L. Carpet Weed. Cultivated ground and waste places in sandy soil; frequent. July-Oct.

CARYOPHYLLACEAE. PINK FAMILY

AGROSTEMMA L. CORN COCKLE

A. GITHAGO L. Cultivated fields; occasional. June, July.

ARENARIA L. SANDWORT

- A. lateriflora L. Blunt-leaved Sandwort. Rocky headland of Lake Champlain, Burlington, (Miss Hitchcock). June, July.
- A. SERPYLLIFOLIA L. Thyme-leaved Sandwort. Dry rocky places; frequent. June-Aug.
- A. stricta Michx. Rock Sandwort. Cliffs and headlands of Lake Champlain; occasional. June, July.

CERASTIUM L. MOUSE-EAR CHICKWEED

- C. arvense L. Field Mouse-ear Chickweed. Dry or rocky places; rare. June, July. Colchester, (Griffin).
- C. nutans Raf. Nodding Chickweed. Moist rich soil; rare. June, July. Colchester, (Griffin).
- C. VULGATUM L. Common Mouse-ear Chickweed. Fields, dooryards, etc.; common. Apr.-Oct.

DIANTHUS L. PINK. CARNATION

D. BARBATUS L. Sweet William. Roadside escape from garden, Essex, (N. F. F.). June, July.

GYPSOPHILA L.

 G_{\star} Muralis L. Low Gypsophila. Spontaneous in garden, Burlington, (Jones); in old field, Colchester, (N. F. F.). July-Sept.

LYCHNIS (Tourn.) L. CAMPION

I. ALBA Mill. White Campion. Waste grounds and roadsides; rare. June-Aug. Burlington, (Grout); weed in meadow, Colchester, (Jones). L. Red Campion. Persistent in old dooryard, Colchester, (N. F. F.); in grass land, Burlington, (Morse). July-Sept.

SAPONARIA L.

- S. officinalis L. Soapwort, Bouncing Bet. Waste places and roadsides; common. July-Oct.
- S. VACCARIA L. Cow-herb. Railroad yards, Burlington, (N. F. F., Ross). June, July.

SILENE L. CATCHFLY. CAMPION

- S. antirrhina L. Sleepy Catchfly. Dry soil; frequent. June-Aug.
- S. Armeria L. Sweet William Catchfly. Spontaneous in garden, Burlington, (N. F. F.). July-Sept.
- S. LATIFOLIA (Mill.) Britten & Rendle. Bladder Campion. Roadsides and fields; occasional. June, July.
- S. NOCTIFLORA L. Night-flowering Catchfly. Cultivated grounds and waste places; occasional. June-Oct.

SPERGULA L. SPURREY

- S. ARVENSIS L. Corn Spurrey. Weed in cultivated grounds; occasional. June-Sept.
- S. Sativa Boenn. Field Spurrey. Adventive at Experiment Farm, Burlington, (Hazen). June, July.

STELLARIA L. CHICKWEED. STARWORT

- S. borealis Bigel. Northern Stitchwort. Shaded or wet places; rare. June-Aug. Burlington, (N. F. F.).
- S. GRAMINEA L. Lesser Stitchwort or Starwort. Roadsides and moist grasslands; frequent. June, July. Var. LATIFOLIA Peterm. Damp grassy place, Williston, (N. F. F.).
- S. longifolia Muhl. Long-leaved Stitchwort. Damp intervals; occasional. June, July.
- S. MEDIA (L.) Cyrill. Common Chickweed. A common weed in moist shaded places. Apr.-Oct.

PORTULACACEAE. PURSLANE FAMILY

CLAYTONIA (Gronov.) L. SPRING BEAUTY

- C. caroliniana Michx. Spring Beauty. Rich open woods; frequent.

 Apr., May.
- C. virginica L. Broad-leaved Spring Beauty. Wooded edge of interval, South Burlington, (Torrey, Jones). Apr., May.

PORTULACA (Tourn.) L. PURSLANE

P. OLERACEA L. Common Purslane, Pusley. Cultivated and waste grounds; common. June-Aug.

NYMPHAEACEAE. WATER LILY FAMILY

BRASENIA Schreb. WATER SHIELD

B. Schreberi Gmel. Ponds; rare. July, Aug. Shelburne, (Jones); Colchester, (Macrae, N. F. F.).

CASTALIA Salisb. WATER LILY. WATER NYMPH

- C. odorata (Ait.) Woodville & Wood. Sweet-Scented White Water Lily, Pond Lily. Ponds and slow streams; common. July, Aug.
- C. tuberosa (Paine) Greene. Tuberous White Water Lily. Tributaries of Lake Champlain; rare. July, Aug. Shelburne, (Pringle); Burlington, (Grout and Tracy).

NYMPHAEA (Tourn.) L. YELLOW POND LILY. SPATTER DOCK

- N. advena Ait. Cow Lily, Large Yellow Pond Lily. Ponds and slow streams; common. June-Aug.
- N. microphylla Pers. Small Yellow Pond Lily. Ponds and slow streams; frequent. June-Aug.
- X ? N. rubrodisca (Morong) Greene. La Plot River, Shelburne, (Pringle). June-Aug.

RANUNCULACEAE. CROWFOOT FAMILY

ACTAEA L. BANEBERRY, COHOSH

- A. alba (L.) Mill. White Baneberry, Doll's Eyes. Rich woods; common. May.
- A. rubra (Ait.) Willd. Red Baneberry. Rich woods; common. May.

ANEMONE (Tourn.) L. ANEMONE

- A. canadensis L. Round-headed Anemone. Moist rocky banks; frequent. May, June.
- A. cylindrica Gray. Long-fruited Anemone. Dry woods and fields; frequent. June, July.
- A. multifida Poir. Cut-leaved Anemone. Ledgy shores; rare. June. Colchester and Burlington, (Robbins, Torrey).
- A. quinquefolia L. Wood Anemone, Wind Flower. Margin of woods; common. May.
- A. riparia Fern. Gravelly and rocky banks; frequent. June, July.
- A. virginiana L. Tall Anemone. Woods and meadows; common. June, July.

AQUILEGIA (Tourn.) L. COLUMBINE

- A. canadensis L. Wild Columbine. Rocky places and dry sandy plains; common. May, June.
- A. VULGARIS L. Garden Columbine. Common in cultivation and a rare roadside escape. June. Colchester, (Jones); Williston, (N. F. F.).

CALTHA (Rupp.) L. MARSH MARIGOLD

C. palustris L. Cowslips. Swamps and wet meadows; common. Apr., May.

CIMICIFUGA L. BUGBANE

C. racemosa (L.) Nutt. Black Snakeroot, Black Cohosh. U. V. M. Herb., Torrey collection. There is no locality on the label, but there is very little doubt that this was collected in Burlington, fifty years or so ago. July.

CLEMATIS L. VIRGIN'S BOWER

- C. verticillaris DC. Purple Clematis. Rocky woods; frequent. May.
- C. virginiana L. Wild Clematis, Virgin's Bower, Traveler's Joy. River banks and moist thickets; common. July, Aug.

COPTIS Salisb. GOLDTHREAD

C. trifolia (L.) Salisb. Canker-root. Wet woods; common. Apr., May.

HEPATICA (Rupp.) Hill. LIVERLEAF. HEPATICA

H. acutiloba DC. Sharp-lobed Liverleaf. Woods; frequent. Apr., May.

H. triloba Chaix. Round-lobed Liverleaf. Warm open woods; common. Apr., May. Interesting variations in leaf and flower characters have been observed.*

HYDRASTIS Ellis. ORANGE-ROOT. YELLOW PUCCOON

H. canadensis L. Golden Seal. Rich woods; rare. May. Shelburne, (Horsford).

RANUNCULUS (Tourn.) L. CROWFOOT. BUTTERCUP

- R. abortivus L. Small-flowered Crowfoot. Shady banks; common. Apr.-June.
- R. abortivus L., var. eucyclus Fern. Shady bank, Burlington, (N. F. F.). May, June.
- R. ACRIS L. Tall Crowfoot or Buttercup. Everywhere as a weed in grass lands. May-Oct. Var. Steveni (Andrz.) Lange. Grass lands; frequent. May-Oct.
- R. aquatilis L., var. capillaceus DC. Common White Water Crowfoot. In slow water; rare. June, July. Colchester, (Griffin).
- R. Bulbosus L. Bulbous Crowfoot or Buttercup. Introduced in field, South Burlington, (Collins). May, June.
- R. circinatus Sibth. Stiff Water Crowfoot. La Plot River, Shelburne, (Torrey). June-Aug.
- R. delphinifolius Torr. Yellow Water Crowfoot. Still water and muddy banks; occasional. May, June. Var. TERRESTRIS (Gray) Farwell. Occasional. This seems to be the form the species takes when rooting out of water or left in the mud of drying ponds.
- R. Flammula L., var. reptans (L.) Mey. Creeping Spearwort. Gravelly or sandy shores; occasional. July, Aug.
- R. pennsylvanicus L. f. Bristly Crowfoot. Wet places; frequent. July, Aug.
- R. recurvatus Poir. Hooked Crowfoot. Moist woods; frequent. May, June.
- R. REPENS L. Creeping Buttercup. Adventive in lawns, Burlington, (Jones, Mrs. Forbes). June, July.
- R. sceleratus L. Cursed Crowfoot. Wet ditches and low grounds; local. June, July. Colchester and Burlington, (N. F. F.).

^{*}See "Vagaries of Hepatica" by F. A. Ross in Torreya, Vol. 3. (Apr., 1903).

R. septentrionalis Poir. Swamp Buttercup. Wet places; frequent. May, June.

THALICTRUM (Tourn.) L. MEADOW RUE

- T. confine Fern. Rocky shore of Lake Champlain, Burlington, (N. F. F.). July-Sept.
- T. dioicum L. Early Meadow Rue. Rocky woodlands; common. Apr., May.
- T. polygamum Muhl. Tall Meadow Rue. Moist meadows; common. June, July.

BERBERIDACEAE. BARBERRY FAMILY

BERBERIS (Tourn.) L. BARBERRY

B. VULGARIS L. Common Barberry. A frequent escape from cultivation. May, June.

CAULOPHYLLUM Michx. Blue Cohosh

C. thalictroides (L.) Michx. Pappoose Root. Rich woods; common. Apr., May.

LAURACEAE. LAUREL FAMILY

BENZOIN Fabric. WILD ALLSPICE. FEVER BUSH

B. aestivale (L.) Nees. Spice Bush, Benjamin Bush. Damp woods. Apr., May. Abundant at the one station, South Burlington, (N. F. F.).

PAPAVERACEAE. POPPY FAMILY

CHELIDONIUM (Tourn.) L. CELANDINE

C. MAJUS L. Waste places; frequent. May-Aug.

PAPAVER (Tourn.) L. POPPY

P. RHOEAS L. Corn Poppy. Persistent in garden, Burlington, (N. F. F.). June-Sept. This is the commonly cultivated Shirley poppy.

SANGUINARIA (Dill.) L. BLOODROOT

S. canadensis L. Borders of rich woods; common. Apr., May.

FUMARIACEAE. FUMITORY FAMILY

ADLUMIA Raf. CLIMBING FUMITORY

A. fungosa (Ait.) Greene. Mountain Fringe, Allegheny Vine. Rich rocky woods; occasional. July-Sept.

CORYDALIS (Dill.) Medic.

- C. aurea Willd. Golden Corydalis. Rocky banks; rare. Apr.-June. Burlington, (Torrey); Colchester, (Griffin).
- C. sempervirens (L.) Pers. Pale or Pink Corydalis. Rocky woodlands; frequent. May-Aug.

DICENTRA Bernh.

- D. canadensis (Goldie) Walp. Squirrel Corn. Rich woods; rare. Apr., May. Shelburne, (Torrey); South Burlington, (N. F. F.).
- D. Cucullaria (L.) Bernh. Dutchman's Breeches. Rich woods; occasional. Apr., May.

FUMARIA (Tourn.) L. FUMITORY

F. OFFICINALIS L. Common Fumitory. Persistent in old garden, Essex, (Jones). June-Aug.

CRUCIFERAE. MUSTARD FAMILY

ARABIS L. ROCK CRESS

- A. brachycarpa (T. & G.) Britton. Purple Rock Cress. Sandy soil, rocky banks, etc.; frequent. June, July.
- A. Drummondii Gray. Rocky places; occasional. May, June.
- A. glabra (L.) Bernh. Tower Mustard. Rocks and fields; occasional.

 June.
- A. hirsuta (L.) Scop. Hairy Rock Cress. Rocky places; frequent. June.
- A. laevigata (Muhl.) Poir. Smooth Rock Cress. Rocky places; occasional. May, June.

BARBAREA R. Br. WINTER CRESS

B. stricta Andrz. Erect-fruited Winter Cress. Roadsides and meadows; common. May, June.

B. VULGARIS R. Br. Common Winter Cress, Yellow Rocket. Low grounds and roadsides; rare. May, June. South Burlington, (N. F. F.).

BERTEROA DC.

B. INCANA (L.) DC. Hoary Alyssum. A weed in lawns and by waysides. First noticed in Burlington, (Miss Towle), 1895, but now becoming frequent. June-Oct.

BRASSICA (Tourn.) L. MUSTARD. TURNIP

- B. ARVENSIS (L.) Ktze. Charlock. A weed in fields; common. June. Oct.
- B. CAMPESTRIS L. Rutabaga. Fields; frequent. June-Oct.
- B. JUNCEA (L.) Cosson, Indian Mustard. Fields; occasional. June-Oct.
- B. NIGRA (L.) Koch. Black Mustard. Waste places; rarely seen. July-Sept. Burlington, (N. F. F.).

CAMELINA Crantz. FALSE FLAX

C. MICROCARPA Andrz. Fields, etc.; occasional. June, July.

CAPSELLA Medic. SHEPHERD'S PURSE

C. Bursa-pastoris (L.) Medic. A common weed. Apr.-Oct.

CARDAMINE (Tourn.) L. BITTER CRESS

- C. parviflora L. Small Bitter Cress. Dry rocky woods; occasional.

 May, June.
- C. pennsylvanica Muhl. Bitter Cress. Cold wet woods and moist ground; common. May, June.

DENTARIA (Tourn.) L. TOOTHWORT. PEPPER-ROOT

- D. diphylla Michx. Two-leaved Toothwort or Pepper-root. Rich moist woods; occasional. May.
- D. laciniata Muhl. Cut-leaved Toothwort or Pepper-root. Rich damp woods; rare. May. South Burlington, (Torrey); Shelburne, (Pringle).
- D. maxima Nutt. Large Toothwort or Pepper-root. Rich moist woods; rare. May. Shelburne, (Pringle); South Burlington, (Perkins); Colchester, (Griffin).

DRABA (Dill.) L. WHITLOW-GRASS

D. arabisans Michx. Twisted Whitlow-grass. Rocky cliffs; rare. May, June. Burlington, (Torrey).

ERYSIMUM (Tourn.) L. TREACLE MUSTARD

E. CHEIRANTHOIDES L. Worm-seed Mustard. Fields and railroad yards; occasional. July-Oct.

LEPIDIUM (Tourn.) L. PEPPERWORT. PEPPERGRASS

- L. APETALUM Willd. Wild Peppergrass. Dry soil, especially along railroads; common. June-Oct.
- L. CAMPESTRE (L.) R. Br. Field or Cow Cress. Fields and roadsides; rare. June. Burlington, (Jones).
- L. VIRGINICUM L. Wild Peppergrass. Dry soil, especially along railroads; common. June-Oct.

LOBULARIA Desy. Sweet Alyssum

L. MARITIMA (L.) Desv. Occasionally spontaneous in gardens and waste places. July-Sept.

RADICULA (Dill.) Hill. WATER CRESS

- R. aquatica (Eat.) Robins. Lake Cress. Lakes and rivers; rare. July, Aug. La Plot River, Shelburne, (Pringle); Malletts Creek, Colchester, (Griffin).
- R. Armoracia (L.) Robins. Horseradish. A frequent escape to moist grounds. May, June.
- R. palustris (L.) Moench. Marsh Cress. Wet places; occasional. June, July. Burlington, (N. F. F.).
- R. palustris (L.) Moench, var. hispida (Desv.) Robins. Hairy Marsh Cress. Wet places; frequent. June, July,
- R_{\bullet} sylvestris (L.) Druce. Yellow Cress. A weed in cultivated ground and wet meadow, South Burlington, (N. F. F.). June.

SISYMBRIUM (Tourn.) L. HEDGE MUSTARD

S. ALTISSIMUM L. Tumble Mustard. First introduced a few years ago, but already becoming a common weed in fields, roadsides, etc. June-Oct.

S. OFFICINALE (L.) Scop., var. Leiocarpum DC. Common Hedge Mustard. A common weed in waste places and cultivated grounds.

June-Oct.

THLASPI (Tourn.) L. PENNY CRESS

T. ARVENSE L. Field Penny Cress, Mithridate Mustard. Waste places; occasional. June. Burlington, (Jones, N. F. F.).

CAPPARIDACEAE. CAPER FAMILY

POLANISIA Raf.

P. graveoleus Raf. Clammy-weed. Sandy shore of Lake Champlain; common. July, Aug.

SARRACENIACEAE. PITCHER-PLANT FAMILY

SARRACENIA (Tourn.) L.

S. purpurea L. Side-saddle Flower, Pitcher-plant, Huntsman's Cup. Sphagnum bogs; frequent. June.

DROSERACEAE. SUNDEW FAMILY

DROSERA L. SUNDEW

- **D. longifolia** L. Long-leaved Sundew. Bogs; rare. July, Aug. Colchester, (Pringle, Griffin).
- D. rotundifolia L. Round-leaved Sundew. Bogs and wet banks; frequent. July, Aug.

CRASSULACEAE. ORPINE FAMILY

PENTHORUM (Gronov.) L. DITCH STONECROP

P. sedoides L. Ditches and swamps; frequent. July-Sept.

SEDUM (Tourn.) L. STONECROP. ORPINE

- S. ACRE L. Mossy Stonecrop. On rocks; occasional. June, July.
- S. PURPUREUM Tausch. (S. triphyllum (Haw.) Gray) Garden Orpine, Live-for-ever. Fields and roadsides; common. Aug., Sept.

S. TELEPHIOIDES Michx. American Orpine. A roadside escape in sandy soil, Colchester, (N. F. F.). Sept.

SAXIFRAGACEAE. SAXIFRAGE FAMILY

CHRYSOSPLENIUM (Tourn.) L. GOLDEN SAXIFRAGE

C. americanum Schwein. Wet shady places; frequent. May, June.

MITELLA (Tourn.) L. MITERWORT. BISHOP'S CAP

- M. diphylla L. Two-leaved Miterwort. Rich woods; common. Apr., May.
- M. nuda L. Naked Miterwort. Deep moist woods, in moss; rare. May, June. South Burlington, (Jones); Burlington, (N. F. F.).

PARNASSIA (Tourn.) L. GRASS OF PARNASSUS

P. caroliniana Michx. Wet banks; rare. Aug. South Burlington, (Torrey, Ross); Burlington, (N. F. F.).

RIBES L. CURRANT. GOOSEBERRY

- R. AUREUM Pursh. (R. odoratum Wend.) Missouri or Buffalo Currant. Occasionally persistent about old building sites. May.
- R. Cynosbati L. Prickly Gooseberry, Dogberry. Rocky woods; common. May.
- R. floridum L'Her. (R. americanum Mill.) Wild Black Currant. Moist rich woods; frequent. May, June.
- R. triste Pall. Swamp Red Currant. Cold swamps; occasional. May.
- R. VULGARE Lam. Garden Red Currant. Common in cultivation and occasional as an escape in fence-rows and waste places. Apr., May.

SAXIFRAGA (Tourn.) L. SAXIFRAGE

- S. pennsylvanica L. Swamp Saxifrage. Bogs; rare. May, June. South Burlington, (Benedict, N. F. F.).
- S. virginiensis Michx. Early Saxifrage. Exposed rocks and dry hill-sides; common. Apr., May.

TIARELLA L. FALSE MITERWORT

T. cordifolia L. Foam-flower. Rich moist woods; common. May, June.

HAMAMELIDACEAE. WITCH-HAZEL FAMILY

HAMAMELIS L. WITCH-HAZEL

H. virginiana L. Low woods; common. Oct.

PLATANACEAE. PLANE TREE FAMILY

PLATANUS (Tourn.) L. SYCAMORE. BUTTONWOOD

P. occidentalis L. Plane Tree. Alluvial banks; frequent. May.

ROSACEAE. ROSE FAMILY

AGRIMONIA (Tourn.) L. AGRIMONY

- A. gryposepala Wallr. Tall Hairy Agrimony. Roadsides and borders of woods; frequent. July-Sept.
- A. striata Michx. Woodland Agrimony. Woods and thickets; occasional. July, Aug.

AMELANCHIER Medic. JUNEBERRY

- A. canadensis (L.) Medic. Shad Bush, Service Berry. Woods; common. Apr., May.
- A. oblongifolia (T. & G.) Roem. Rock Shad Bush. Moist woods and rocky uplands; frequent. May.
- A. spicata (Lam.) Koch. Shore Shad Bush. Banks of streams and sandy soil; frequent. May.

CRATAEGUS (Tourn.) L. HAWTHORN. WHITE THORN

- C. alnorum Sarg. Burlington, (Eggleston). Fl. May; fr. Sept.
- C. apposita Sarg. Burlington, (Edson). Fl. May, June; fr. Oct.
- C. macracantha Lodd. Burlington and South Burlington, (Edson). Fl. May; fr. Sept.
- C. macrosperma Ashe. Burlington, (Edson); Essex, (Jones). Fl. May; fr. Sept. Var. MATURA (Sarg.) Eggl. Burlington, (Eggleston). Fl. May; fr. Aug., Sept.
- C. OXYACANTHA L. (C. monogyna Jacq.) English Hawthorn. Naturalized in ravines about Burlington, (Jones). Fl. May; fr. Sept.

- ('. pruinosa (Wendl.) Koch. Burlington, (Edson). Fl. May; fr. Oct., Nov. Var. PHILADELPHICA (Sarg.) Eggl. Burlington, (Eggleston).
- C. punctata Jacq. South Burlington, (Jones); Burlington, (Edson).
 Fl. May, June; fr. Oct.
- C. rotundifolia Moench. Burlington, (Edson). Fl. May; fr. Sept.
- C. silvicola Beadle, var. Beckwithae (Sarg.) Eggl. Burlington, (Eggleston). Fl. May; fr. Oct.

DALIBARDA Kalm.

D. repens L. Cool moist woods; frequent. July.

FILIPENDULA (Tourn.) Hill.

F. VULGARIS Moench. Dropwort. A garden escape well established by roadside, Burlington, (N. F. F.). Station destroyed by street improvements in 1908. June.

FRAGARIA (Tourn.) L. STRAWBERRY

- F. vesca L., var. americana Porter. Wood Strawberry. Woods; common. Apr.-June.
- F. virginiana Duch. Wild Strawberry. Fields; common. Apr., May.

GEUM L. AVENS

- G. canadense Jacq. White Avens. Borders of woods; common. July, Aug.
- G. rivale L. Water or Purple Avens. Bogs or wet meadows; occasional. May, June.
- G. strictum Ait. Yellow Avens. Moist meadows; common. July, Aug.
- G. virginianum L. Rough Avens. Low grounds; rare. July. Burlington, (N. F. F.).

POTENTILLA L. CINQUEFOIL. FIVE-FINGER

- P. Anserina L. Silver Weed. Shores of Lake Champlain; common. June-Aug.
- P. anserina L., var. concolor Ser. (Var. sericea Hayne) Shore of Lake Champlain, Burlington, (N. F. F.). Aug.
- P. argentea L. Silvery Cinquefoil. Dry barren fields; common. June-Sept.

- P. arguta Pursh. Tall Cinquefoil. Rocky hills; rare. June, July. South Burlington, (Jones); Colchester, (Griffin).
- P. canadensis L., var. simplex (Michx.) T. & G. Five-finger. Dry sandy soil; common. May-July.
- P. fruticosa L. Shrubby Cinquefoil. Frequent in the boggy meadows in the vicinity of Winooski Gorge. May-Sept.
- P. monspeliensis L. Rough Cinquefoil. Dry soil; common. May-Aug.
- P. palustris (L.) Scop. Marsh Cinquefoil. Cool bogs; frequent. June, July.
- P. RECTA L. Adventive in lawn, Burlington, (Hazen). Aug.

PRUNUS (Tourn.) L. PLUM. CHERRY

- P. cuneata Raf. Appalachian Cherry. Sandy shores of Lake Champlain and on the sand plains; frequent. May.
- P. nigra Ait. Wild or Canada Plum. Woods and fencerows; frequent. Apr., May.
- P. pennsylvanica L. f. Wild Red, Bird, Fire, or Pin Cherry. Dry woods and clearings; common. Apr., May.
- P. pumila L. Sand Cherry. Rocky shore of Winooski River; rare. May.
- P. serotina Ehrh. Wild Black or Rum Cherry. Woodlands; frequent. May, June.
- P. virginiana L. Choke Cherry. Fencerows, woods and banks; common. May, June.

PYRUS (Tourn.) L.

- P. americana (Marsh.) DC. American Mountain Ash. Wet woods; occasional. May, June.
- P. arbutifolia (L.) L. f. Red Chokeberry. Swamp, Burlington, (Rand). May, June.
- P. arbutifolia (L.) L. f., var. atropurpurea (Britton) Robins. Purple Chokeberry. Swamp, South Burlington, (N. F. F.). May, June.
- P. AUCUPARIA (L.) Ehrh. European Mountain Ash, Rowan Tree. Naturalized and frequent about Burlington. May.
- P. Malus L. Apple. Fencerows and roadsides; frequent. May.
- P. melanocarpa (Michx.) Willd. Black Chokeberry. Moist woods, and also rocky uplands; common. May, June.

ROSA (Tourn.) L. Rose

- R. acicularis Lindl., var. Bourgeauiana Crepin. Prickly Rose. Ledges, rocky woods, etc.; rare. June, July. Burlington, (Brainerd).
- R. blanda Ait. Smooth Rose. Roadsides and ledges; frequent. June, July.
- R. carolina L. Swamp Rose. Borders of swamps and streams; occasional. June, July.
- R. CINNAMOMEA L. Cinnamon Rose. Once much cultivated and now established by roadsides, etc.; occasional. June, July. South Burlington, (Jones); Colchester, (N. F. F.).
- R. humilis Marsh. Pasture Rose. Rocky slopes; frequent. June, July.
- R. RUBIGINOSA L. Sweetbrier, Eglantine. Old pastures; frequent. June, July.

RUBUS (Tourn.) L. BRAMBLE

- R. allegheniensis Porter. High-bush Blackberry. Dry open thickets and recent clearings; frequent. May, June.
- R. hispidus L. Running Swamp Blackberry, Swamp Dewberry. Moist woods; frequent. June, July.
- R. idaeus L., var. aculeatissimus (Mey.) Regel & Tiling. (R. idaeus L., var. canadensis Rich.) Wild Red Raspberry. Thickets and clearings; common. May, June.
- ? R. neglectus Peck. Purple Raspberry. Rocky woods, gravelly banks, etc.; occasional. May, June.
- R. occidentalis L. Black Raspberry, Thimbleberry. Copses, fencerows, etc.; frequent. May, June. Forma Pallidus (Bailey) Robins. Fencerow, Burlington, (Sprague).
- R. odoratus L. Purple Flowering Raspberry. Moist shaded hillsides; common. July, Aug.
- R. recurvans Blanch. Recurved Blackberry. Open soil, thickets, etc.; occasional. June.
- R. triilorus Richards. (R. pubescens Raf.) Dwarf or Running Raspberry, Red Dewberry. Wet woods; frequent. May, June.
- R. villosus Ait. Black Dewberry. Dry open places; occasional. May, June.

SANGUISORBA (Rupp.) L. BURNET

S. MINOR Scop. Garden Burnet. Established in grass lands, Shelburne, (N. F. F.). June.

SORBARIA A. Br.

S. SORBIFOLIA (L.) A. Br. Ash-leaved Spiraea. An occasional escape to copses and roadsides, where it is well established. June.

SPIRAEA (Tourn.) L.

- S. latifolia (Ait.) Borkh. Meadow-sweet, Queen of the Meadow. Wet meadows and rocky pastures; common. July, Aug.
- S. tomentosa L. Hardhack, Steeple Bush. Low grounds; common. July, Aug.

WALDSTEINIA Willd.

W. fragarioides (Michx.) Tratt. Barren Strawberry. Wooded hill-sides; frequent. Apr., May.

LEGUMINOSAE. PULSE FAMILY

AMPHICARPA Ell. HOG PEANUT

- A. monoica (L.) Ell. Rich damp woodlands; common. July-Sept.

 APIOS (Boerh.) Ludwig. Groundnut. Wild Bean
- A. tuberosa Moench. Low moist ground; frequent. July-Sept.

ASTRAGALUS (Tourn.) L. MILK VETCH

- A. canadensis L. Shores of Lake Champlain; rare. July, Aug. South Burlington, (Torrey); Colchester, (Griffin).
- A. Rebbinsii (Oakes) Gray. On limestone rocks, Winooski Gorge, (Robbins). This, the only known station in Vermont, was destroyed in 1894 when the new dam flooded the rocks.

CICER L.

C. ARIETINUM L. Chick-pea. Adventive on railway embankment, Colchester, (N. F. F.). Sept., Oct.

DESMODIUM Desv. TICK TREFOIL

D. canadense (L.) DC. Showy Tick Trefoil. Open woods and banks of streams; frequent. July, Aug.

- D. Dillenii Darl. Open woodlands; occasional. July, Aug.
- D. grandiflorum (Walt.) DC. Rich woods; common. July, Aug.
- D. nudiflorum (L.) DC. Dry woods; frequent. July, Aug.

LATHYRUS (Tourn.) L. VETCHLING. EVERLASTING PEA

- L. maritimus (L.) Bigel. Beach Pea. Shores of Lake Champlain; occasional. June-Aug. A white-flowered form occurs.
- L. ochroleucus Hook. Cream-colored Vetchling. Headlands of Lake Champlain; rare. June, July. South Burlington, (Torrey); Colchester, (N. F. F.).
- L. palustris L. Marsh Pea. Shores of Lake Champlain; occasional. July.

LESPEDEZA Michx. BUSH CLOVER

L. capitata Michx. Sandy soil; occasional. Aug.

LUPINUS (Tourn.) L. LUPINE

L. PERENNIS L. Wild Lupine. Sand plains; frequent. May, June.

MEDICAGO (Tourn.) L. MEDICK

- M. ARABICA (L.) Huds. Spotted Medick. Adventive at Experiment Farm, Burlington, (Jones). June-Aug.
- M. HISPIDA Gaertn. Bur Clover. Adventive at Experiment Farm, Burlington, (Jones). June.
- M. LUPULINA L. Black Medick, Nonesuch. Roadsides and fields; common. May-Sept.
- M. SATIVA L. Lucerne, Alfalfa. Often cultivated and occasionally spontaneous. June, July.

MELILOTUS (Tourn.) Hill. SWEET CLOVER. MELILOT

- M. ALBA Desr. White Melilot or Sweet Clover. Waste grounds and along railroads; frequent. June-Oct.
- M. officinalis (L.) Lam. Yellow Melilot or Sweet Clover. Roadsides and waste places; occasional. June-Aug.

ONOBRYCHIS L. SAINFOIN

O. SATIVA Lam. Adventive at Experiment Farm, Burlington, (Jones).
June.

ROBINIA L. LOCUST

- R. PSEUDO-ACACIA L. Common Locust, False Acacia. Frequent in cultivation and thoroughly naturalized. June.
- R. viscosa Vent. Clammy Locust. An established roadside escape, South Burlington and Colchester, (N. F. F.). June.

TRIFOLIUM (Tourn.) L. CLOVER. TREFOIL

- T. AGRARIUM L. Yellow or Hop Clover. Sandy soil; frequent. May-Aug.
- T. ARVENSE L. Rabbit-foot or Stone Clover. Dry sandy soil; common. July-Sept.
- T. HYBRIDUM L. Alsike Clover. Fields and roadsides; common. May-Aug.
- T. MEDIUM L. Zigzag Clover. Adventive at Experiment Farm, Burlington, (Jones). July, Aug.
- T. PRATENSE L. Red Clover. Fields and roadsides; common. May-Sept.
- T. PROCUMBENS L. Low Hop Clover. Railway embankment and lumber yards; rare. June, July. Burlington, (Kirk, N. F. F.).
- T. repens L. White Clover. Fields and roadsides; common. May, June.

VICIA (Tourn.) L. VETCH. TARE

- V. ANGUSTIFOLIA Reich. One form occurs. Var. SEGETALIS (Thuill.)

 Koch. Common Vetch. Roadsides, waste places, etc.; common.

 June-Aug. A white-flowered form occurs on rocky bluff and beach of Lake Champlain, Burlington, (N. F. F.).
- V. CRACCA L. Blue Vetch. Meadows and roadsides; frequent. June-Aug.
- V. TETRASPERMA (L.) Moench. Slender Vetch. In cultivated field, Burlington, (Ross). June, July.

LINACEAE. FLAX FAMILY

LINUM (Tourn.) L. FLAX

L. USITATISSIMUM L. Common Flax. Occasionally adventive. July-Sept.

OXALIDACEAE. WOOD SORREL FAMILY

OXALIS L. WOOD SORREL

- (). Acetosella L. Common Wood Sorrel. Cool moist woods; occasional. June.
- corniculata L. Lady's Sorrel. Fields and gardens; common. June-Sept.
- (). stricta L. Yellow Wood Sorrel. Dry or sandy soil; occasional.

 June-Sept.

GERANIACEAE. GERANIUM FAMILY

ERODIUM L'Her. STORKSBILL

- E. Botrys Bertol. On wool waste, bank of Winooski River, Burlington, (Brainerd). Sept., Oct.
- F. CICUTARIUM (L.) L'Her. Storksbill. Old garden, Burlington, (Mrs. Grout); Experiment Farm, Burlington, (Jones). July, Aug.
- F. MOSCHATUM (L.) L'Her. Musk Storksbill. On wool waste, bank of Winooski River, Burlington, (Brainerd). Sept., Oct.

GERANIUM (Tourn.) L. CRANESBILL

- G. Bicknellii Britton. Dry rocky places; frequent. May, June.
- G. maculatum L. Wild Cranesbill. Open woods and fields; occasional. May, June.
- G. Robertianum L. Herb Robert. Rocky woods and ravines; common. May-July.

RUTACEAE. RUE FAMILY

ZANTHOXYLUM L. PRICKLY ASH

Z. americanum Mill. Northern Prickly Ash, Toothache Tree. Rocky woods and river banks; occasional. Apr., May.

POLYGALACEAE. MILKWORT FAMILY

POLYGALA (Tourn.) L. MILKWORT

- P. paucifolia Willd. Fringed Polygala, Flowering Wintergreen. Woods, in light soil; common. May. A white-flowered form occurs, Burlington, (Bigelow).
- P. polygama Walt. Racemed Milkwort. Dry sandy soil; occasional. June, July.
- P. sanguinea L. Field or Purple Milkwort. Moist sandy soil; rare. July-Sept. Burlington, (N. F. F.).
- P. Senega L. Seneca Snakeroot. Rocky soil; frequent. May, June.
- P. verticillata L., var. ambigua (Nutt.) Wood. Loose-spiked Milkwort. Dry sandy soil; rare. Aug. Burlington, (Howe).

EUPHORBIACEAE. SPURGE FAMILY

ACALYPHA L. THREE-SEEDED MERCURY

A. virginica L. Moist hollows in fields; frequent. July-Sept.

EUPHORBIA L. SPURGE

- E. COROLLATA L. Flowering Spurge. Established in sandy field, Burlington, (N. F. F.). July, Aug.
- E. CYPARISSIAS L. Cypress Spurge, Graveyard Moss. A frequent escape from cultivation about old houses and by roadsides. May-Sept.
- E. HELIOSCOPIA L. Wartweed. Gravelly shores and along Rutland R. R.; frequent. Aug., Sept.
- E. hirsuta (Torr.) Wiegand. Dry sandy soil; common. July-Sept.
- E. maculata L. Milk Purslane. Open places, roadsides, etc.; frequent. July-Sept.
- E. PLATYPHYLLA L. Shore of Lake Champlain, Shelburne, (Pringle, N. F. F.). June-Aug.

CALLITRICHACEAE. WATER STARWORT FAMILY

CALLITRICHE L. WATER STARWORT

C. palustris L. Quiet water; occasional. July, Aug.

LIMNANTHACEAE. FALSE MERMAID FAMILY

FLOERKEA Willd. FALSE MERMAID

F. proserpinacoides Willd. Bank of La Plot River, Shelburne, (Pringle). May.

ANACARDIACEAE. CASHEW FAMILY

RHUS L. SUMACH

- R, copallina L. Dwarf Sumach. Rocky hills; rare. July. Colchester, (Jones, Griffin).
- R. glabra L. Smooth Sumach. Sandy soil; rare. July. Colchester, (N. F. F., Mrs. Walker).
- R. Toxicodendron L. Poison Ivy, Poison Oak. Banks and moist thickets; common. June. Var. RADICANS (L.) Torr. Common. June.
- R. typhina L. Staghorn Sumach. Hillsides; common. June, July.
- R. Vernix L. Poison Sumach, Elder or Dogwood. Swamps; rare. June. Colchester, (Grout, Jones).

AQUIFOLIACEAE. HOLLY FAMILY

ILEX L. HOLLY

- I. verticillata (L.) Gray. Winterberry, Black Alder. Wet thickets; common. June, July.
- I. verticillata (L.) Gray, var. padifolia (Willd.) T. & G. Colchester, (Griffin). June, July.
- I. verticillata (L.) Gray, var. tenuifolia (Torr.) Wats. Mouth of Winooski River, (Eggleston). June, July.

NEMOPANTHUS Raf. MOUNTAIN HOLLY

N. mucronata (L.) Trel. Cool damp woods and swamps; frequent.

May.

CELASTRACEAE. STAFF TREE FAMILY

CELASTRUS L. STAFF TREE. SHRUBBY BITTER-SWEET

C. seandens L. Waxwork, Climbing Bitter-sweet. Moist banks and thickets; common. June.

STAPHYLEACEAE. BLADDER NUT FAMILY

STAPHYLEA L. BLADDER NUT

S. trifolia L. American Bladder Nut. Moist woods and thickets; occasional. May, June.

ACERACEAE. MAPLE FAMILY

ACER (Tourn.) L. MAPLE

- A. Negundo L. Box Elder, Ash-leaved Maple. Frequent on banks of the Winooski River and common in cultivation about Burlington. Apr., May.
- A. pennsylvanicum L. Striped Maple, Moosewood. Rich woods; common. May.
- A. rubrum L. Red or Swamp Maple. Moist woods and swamps; common. Apr., May.
- A. saccharinum L. White or Silver Maple. River banks; frequent. Apr.
- A. saccharum Marsh. Sugar or Rock Maple. Rich woods; common. Apr., May.
- A. saccharum Marsh., var. nigrum (Michx. f.) Britton. Black Sugar Maple. Occasionally planted as a shade tree in Burlington and vicinity, and search will probably reveal it in the moist woodlands bordering the Winooski River or Lake Champlain. May.
- A. spicatum Lam. Mountain Maple. Cool ravines; common. May, June.

SAPINDACEAE. SOAPBERRY FAMILY

AESCULUS L. HORSE-CHESTNUT. BUCKEYE

A. HIPPOCASTANUM L. Common Horse-chestnut. Commonly planted and occasionally spontaneous. May, June.

BALSAMINACEAE. TOUCH-ME-NOT FAMILY

IMPATIENS (Rivin.) L. BALSAM. JEWELWEED

- I. biflora Walt. Spotted Touch-me-not. Moist shady places; common. June-Sept. A form with reddish flowers occurs, Colchester, (N. F. F.).
- I. pallida Nutt. Pale Touch-me-not. Cool moist shady places; occasional. July-Sept.

RHAMNACEAE. BUCKTHORN FAMILY

CEANOTHUS L. RED-ROOT

- C. americanus L. New Jersey Tea. Dry woodlands; common. July.
- C. ovatus Desf. Smaller Red-root. Sandy banks of Lake Champlain, Burlington, (Torrey). June.

RHAMNUS (Tourn.) L. BUCKTHORN

- R. aluifolia L'Her. Alder-leaved Buckthorn. Swamps; occasional.

 June.
- R. CATHARTICA L. Common Buckthorn. Common in cultivation and frequently naturalized in woods and fields. May, June.

VITACEAE. VINE FAMILY

PSEDERA Neck. VIRGINIA CREEPER. WOODBINE

- P. quinquefolia (L.) Greene, var. hirsuta (Donn) Rehder. Disk Creeper. Clinging closely to limestone ledges, Burlington, (Jones). June.
- P. vitaceae (Knerr) Greene. Tendril Creeper. Moist woods, alluvial thickets, etc.; common. June.

VITIS (Tourn.) L. GRAPE

- V. bicolor Le Conte. Summer Grape. Deep gravelly soil and base of cliffs; occasional. June.
- V. vnlpina L. River-bank or Frost Grape. River banks; common.

 June.

TILIACEAE. LINDEN FAMILY

TILIA (Tourn.) L. LINDEN. BASSWOOD

T. americana L. Basswood. Woodlands; common. June, July.

MALVACEAE. MALLOW FAMILY

ABUTILON (Tourn.) Mill. INDIAN MALLOW

A. THEOPHRASTI Medic. Velvet Leaf. Occasional by waysides and as a garden weed. Aug.-Oct. Williston, (Bates); Burlington, (Jones, N. F. F.); Colchester, (Mrs. Walker).

HIBISCUS L. Rose Mallow

H. TRIONUM L. Flower-of-an-hour, Bladder Ketmia. Occasionally adventive in gardens and waste places. July, Aug. Burlington, (Jones, N. F. F., Mrs. Walker).

MALVA (Tourn.) L. MALLOW

- M. Alcea L. European Mallow. A garden escape, Burlington, (N. F. F.). June-Sept.
- M. Moschata L. Musk Mallow. Roadsides and waste places; occasional. June, July.
- M. ROTUNDIFOLIA L. Common Mallow, Cheeses. Waste places and cultivated grounds; common. May-Oct.
- M. VERTICILLATA L. Whorled Mallow. Occasional as a weed in gardens and waste places. Burlington, (Jones). July, Aug.

HYPERICACEAE. ST. JOHN'S=WORT FAMILY

HYPERICUM (Tourn.) L. St. John's-wort

- H. Ascyron L. Great St. John's-wort. Bank of Winooski River near High Bridge, (Perkins, Grout). July, Aug.
- H. canadense L. Wet places; rare. July, Aug. Colchester, (Torrey, N. F. F.).
- H. ellipticum Hook. Pale St. John's-wort. Wet places; frequent. July, Aug.

- H. mutilum L. Small-flowered St. John's-wort. Low grounds; common. July-Sept.
- H. PERFORATUM L. Common St. John's-wort. Fields and roadsides; common. June-Sept.
- H. punctatum Lam. Spotted St. John's-wort. Damp places; frequent. July, Aug.
- H. virginicum L. Marsh St. John's-wort. Swamps; frequent. July, Aug.

CISTACEAE. ROCKROSE FAMILY

HELIANTHEMUM (Tourn.) Mill. ROCKROSE

H. majus BSP. Frostweed. Sterile soil; frequent. June-Aug.

HUDSONIA L

H. tomentosa Nutt. One form occurs. Var. INTERMEDIA Peck. Beach Heather, Sandy beaches of Lake Champlain; occasional. May, June.

LECHEA (Kalm) L. PINWEED

L. intermedia Leggett. Dry sterile soil; frequent. July, Aug.

VIOLACEAE. VIOLET FAMILY

VIOLA (Tourn.) L. VIOLET. HEART'S-EASE

- V. affinis Le Conte. Moist thickets and boggy meadows. May. Colchester, (Brainerd).
- V. arenaria DC. Sand Violet. Sand plains; common. May.
- V. blanda Willd. Sweet White Violet. Cool rich woods; common. May.
- V. canadensis L. Canada Violet. Woods; common. May, June. Occasional blossoms occur all summer.
- V. conspersa Reich. Dog Violet. Low shady grounds; common. Apr., May.
- V. cucullata Ait. Marsh Blue Violet. Wet places; common. May, June.

- V. fimbriatula Sm. Sandy fields and dry hillsides; common. Apr., May. A white-flowered form occurs, Colchester, (Jones, N. F. F.). A form with the basal lobes coarsely toothed occurs, shore of Burlington Bay (Brainerd).
- V. incognita Brain. Low moist woodlands; rare. May. Burlington, (Grout).
- V. lanceolata L. Lance-leaved Violet. Moist meadow, Williston, (Bates). May, June.
- V. latiuscula Greene. Dry open woods, in light soil; occasional. May. Burlington and Colchester, (Brainerd); Williston, (Ross).
- V. nephrophylla Greene. Cold mossy bogs and borders of streams. Apr., May. Colchester, (Brainerd); South Burlington, (Jones).
- V. pallens (Banks) Brain. White Violet. Low wet grounds; common. May.
- V. pubescens Ait. Downy Yellow Violet. Rich woods; common. Apr., May.
- V. renifolia Gray. White Violet. Wet mossy woods and swamps; occasional. May.
- V. rostrata Pursh. Long-spurred Violet. Rich woods; frequent. Apr., May.
- V. rotundifolia Michx. Round-leaved or Early Yellow Violet. Cool moist woods; frequent. Apr., May.
- V. scabriuscula Schwein. Smooth Yellow Violet. Rich moist woods; frequent. May.
- Y. sororia Willd. Woolly Blue Violet. Moist meadows, alluvial woods, etc.; frequent. May, June. A white-flowered form occurs, Williston, (Ross).
- V. TRICOLOR L. Pansy, Heart's-ease. Persistent in old gardens. Apr.-Oct.

THYMELAEACEAE. MEZEREUM FAMILY

DAPHNE L. MEZEREUM

D. Mezereum L. Frequent in gardens and escaped to rocky woods, Burlington, (Jones). Apr.

DIRCA L. LEATHERWOOD, MOOSEWOOD

D. palustris L. Wicopy. Moist open woods; frequent. Apr., May.

ELAEAGNACEAE. OLEASTER FAMILY

SHEPHERDIA Nutt.

S. canadensis (L.) Nutt. Canadian Buffalo-berry. Rocky bluffs; occasional. Apr., May.

LYTHRACEAE. LOOSESTRIFE FAMILY

DECODON Gmel. SWAMP LOOSESTRIFE

D. verticillatus (L.) Ell. Water Willow. Swampy grounds; occasional. July, Aug.

LYTHRUM L. LOOSESTRIFE

L. ALATUM Pursh. Waste place, Burlington, (N. F. F.). June, July.

ONAGRACEAE. EVENING PRIMROSE FAMILY

CIRCAEA (Tourn.) L. ENCHANTER'S NIGHTSHADE

- C. alpina L. Smaller Enchanter's Nightshade. Deep woods; common. July, Aug.
- C. intermedia Ehrh. Burlington, (Jones); Colchester, (Griffin). July, Aug.
- C. lutetiana L. Enchanter's Nightshade. Woods; frequent. July, Aug.

EPILOBIUM (Dill.) L. WILLOW-HERB

E. adenocaulon Haussk. Northern Willow-herb. Moist places; frequent. July, Aug.

- E. angustifolium L. Great Willow-herb, Fireweed. Common, especially in newly cleared lands. July-Sept. A white-flowered form occurs, Colchester, (Perkins).
- E. coloratum Muhl. Purple-leaved Willow-herb. Wet places; common. July, Aug.
- E. densum Raf. Open low grounds; occasional. July, Aug.
- E. molle Torr. Bogs; rare. Aug., Sept. Colchester, (Torrey, N. F. F.).

LUDVIGIA L. FALSE LOOSESTRIFE

I. palustris (L.) Ell. Water Purslane. Ditches and swamps; common. June-Aug.

OENOTHERA L. EVENING PRIMROSE

- 0. biennis L. Common Evening Primrose. Dry soil; common. July-Sept.
- **0.** cruciata Nutt. Small-flowered Evening Primrose. Sandy or gravelly soil; rare. July-Sept. Burlington, (Ross); Colchester, (Ross, Griffin).
- 0. LACINIATA Hill. Sinuate-leaved Evening Primrose. Adventive at Fort Ethan Allen, (Grout). Sept.
- **0. muricata** L. Sandy soil, Burlington, (N. F. F.). July-Sept. Further search will probably show this to be frequent.
- OAKESIANUS Robbins. Adventive by roadside, Burlington, (Hazen). July-Sept.
- 0. pumila L. Small Sundrops. Fields; common. June-Aug.
- 0. SERRULATA Nutt. Adventive on railroad embankment, Burlington, (N. F. F.). July-Sept.

HALORAGIDACEAE. WATER MILFOIL FAMILY

MYRIOPHYLLUM (Vaill.) L. WATER MILFOIL

M. spicatum L. Shelburne Pond, (Jones); Malletts Bay, (Griffin, N. F. F.). Aug.

ARALIACEAE. GINSENG FAMILY

ARALIA (Tourn.) L.

- A. hispida Vent. Bristly Sarsaparilla, Wild Elder. Rocky or sandy woodlands; frequent. June, July.
- A. nudicaulis L. Wild Sarsaparilla. Moist woodlands; common. May, June.
- A. racemosa L. Spikenard. Rich woodlands; frequent. July, Aug.

PANAX L. GINSENG

- P. quinquefolium L. Ginseng. Rich and cool woods; rare. July. Burlington, (Ross); Colchester, (Griffin).
- P. trifolium L. Dwarf Ginseng, Ground-nut. Moist woods; common. May, June.

UMBELLIFERAE. PARSLEY FAMILY

ANGELICA L. ANGELICA

A. atropurpurea L. Masterwort. Swamps and moist grounds; frequent. June, July.

CARUM (Rupp.) L. CARAWAY

C. CARVI L. Caraway. About dwellings and by roadsides; frequent. May, June.

CICUTA L. WATER HEMLOCK

- C. bulbifera L. Water Hemlock. Swamps; occasional. July.
- C. maculata L. Spotted Cowbane, Musquash Root. Low grounds; frequent. July.

CONIUM L. POISON HEMLOCK

C. MACULATUM L. Waste places; occasional. July. Burlington, (Grout); Colchester, (Jones).

CRYPTOTAENIA DC. HONEWORT

C. canadensis (L.) DC. Moist woods; frequent. June.

DAUCUS (Tourn.) L. CARROT

D. CAROTA L. Wild Carrot, Queen Anne's Lace. A common and pernicious weed in meadows, fields, etc. July-Oct. The pink-flowered form is occasional.

ERYNGIUM (Tourn.) L. ERYNGO

E. VIRGATUM Lam. Roadside escape from garden, Burlington, (Jones).

Aug.

HERACLEUM L. COW PARSNIP

H. lanatum Michx. Masterwort. Wet grounds; occasional. June, July.

HYDROCOTYLE (Tourn.) L. WATER PENNYWORT

H. americana L. Wet places; common. June-Aug.

LEVISTICUM (Rivin.) Hill. LOVAGE

L. officinale (L.) Koch. Roadside escape from gardens; rare. June, July. Shelburne, (N. F. F.).

OSMORHIZA Raf. SWEET CICELY

- Claytoni (Michx.) Clarke. Woolly Sweet Cicely. Rich woods; frequent. May, June.
- longistylis (Torr.) DC. Smoother Sweet Cicely. Rich moist woods; occasional. May, June.

PASTINACA L. PARSNIP

P. SATIVA L. Wild or Poison Parsnip. Roadsides and waste places; common. June-Sept.

SANICULA (Tourn.) L. SANICLE. BLACK SNAKEROOT

- S. canadensis L. Short-styled Snakeroot. Shaded ledges; frequent. July.
- S. gregaria Bick. Clustered Snakeroot. Moist rich soil; rare. July, Aug. South Burlington, (Orton); Colchester, (N. F. F.).
- S. marilandica L. Black Snakeroot. Moist woods; common. June, July.
- S. trifoliata Bick. Large-fruited Snakeroot. Shelburne, (Boott). Specimen in Gray Herb. July.

SIUM (Tourn.) L. WATER PARSNIP

S. cicutaefolium Schrank. Wet places; frequent. Aug., Sept.

TAENIDIA Drude.

T. integerrima (L.) Drude. Yellow Pimpernel. Rocky places; occasional. June, July.

ZIZIA Koch.

Z. aurea (L.) Koch. Golden Alexanders, Early or Golden Meadow Parsnip. Moist meadows; common. May, June.

CORNACEAE. DOGWOOD FAMILY

CORNUS (Tourn.) L. CORNEL. DOGWOOD

- C. alternifolia L. f. Alternate-leaved Dogwood. Open woods; frequent. May, June.
- C. Amomum Mill. Silky Cornel or Dogwood, Kinnikinnik. Low woods and banks; frequent. June, July.
- C. canadensis L. Dwarf Cornel, Bunchberry. Cold damp woods; frequent. May, June.
- C. circinata L'Her. Round-leaved Cornel or Dogwood. Rocky woodlands; frequent. May, June.
- C. paniculata L'Her. Panicled Dogwood. Rich thickets and banks; frequent. June, July.
- C. stolonifera Michx. Red-osier Dogwood. Wet places; common. May, June. Occasional blossoms occur all summer.

NYSSA L. TUPELO. PEPPERIDGE. SOUR GUM

N. sylvatica Marsh. Black Gum. Rich soil, generally swampy; rare. May. Burlington, (N. F. F., Jones); Colchester, (Griffin).

ERICACEAE. HEATH FAMILY

ANDROMEDA L.

A. glaucophylla Link. Bog Rosemary. Bogs; frequent. May, June.

ARCTOSTAPHYLOS Adans. BEARBERRY

A. Uva-ursi (L.) Spreng. Red Bearberry. Rocky bluffs and sand plains; rare. May. Burlington, (Perkins); South Burlington, (N. F. F.).

CHAMAEDAPHNE Moench. LEATHER LEAF. CASSANDRA

C. calyculata (L.) Moench. Bogs; frequent. May.

CHIMAPHILA Pursh. PIPSISSEWA

C. umbellata (L.) Nutt. Prince's Pine, Pipsissewa. Dry woods; frequent. July.

CHIOGENES Salisb. CREEPING SNOWBERRY

C. hispidula (L.) T. & G. Moxie Plum, Capillaire. Peat bogs and mossy woods; rare. May. South Burlington; Colchester, (Griffin).

EPIGAEA L. GROUND LAUREL. TRAILING ARBUTUS

E. repens L. Mayflower. Sandy woodlands; common. Apr., May.

GAULTHERIA (Kalm) L. AROMATIC WINTERGREEN

G. procumbens L. Teaberry, Checkerberry. Cool woods; common. July, Aug.

GAYLUSSACIA HBK. HUCKLEBERRY

G. baccata (Wang.) Koch. Black Huckleberry. Sterile fields and thickets; common. May, June.

KALMIA L. LAUREL

- K. angustifolia L. Sheep Laurel, Lambkill, Wicky. Sterile fields and peat bogs; common. June, July.
- K. polifolia Wang. Pale Laurel. Cold peat bogs; frequent. May, June.

LEDUM (Rupp.) L. LABRADOR TEA

L. groenlandicum Oeder. Cold bogs; frequent. June.

MONESES Salisb. ONE-FLOWERED PYROLA

M. uniflora (L.) Gray. Deep cold woods; rare. June. Burlington, (Torrey, N. F. F.).

MONOTROPA L. INDIAN PIPE. PINESAP

- M. Hypopitys L. Pinesap, False Beech Drops. Rich dry woods; occasional. July-Sept.
- M. uniflora L. Indian Pipe, Corpse Plant. Dark rich woods; common. June-Aug.

PTEROSPORA Nutt. PINE DROPS

P. andromedea Nutt. Pine woods; rare. July, Aug. Williston, (Pringle); Colchester, (Bates, Ross).

PYROLA (Tourn.) L. WINTERGREEN. SHIN LEAF

- P. americana Sweet. Round-leaved Wintergreen. Open or sandy woods; occasional. July.
- P. asarifolia Michx., var. incarnata (Fisch.) Fern. Bog Wintergreen. Sphagnum bogs; rare. July. South Burlington, (N. F. F.).
- P. chlorantha Sw. Greenish-flowered Wintergreen. Open woods; frequent. July.
- P. elliptica Nutt. Shin Leaf. Rich woods; common. July.
- P. secunda L. One-sided Wintergreen. Rich woods; frequent. July.

RHODODENDRON L.

- R. canadense (L.) BSP. Rhodora. Swamps; occasional. May, June.
- R. canescens (Michx.) Don. Swamp or June Pink, Pink Azalea. One bush seen, but unfortunately no specimen taken. May, June. Burlington, (N. F. F.).

VACCINIUM L. BLUEBERRY. CRANBERRY

- V. atrococcum (Gray) Heller. Black High Blueberry. Swamp, South Burlington, (Jones). May.
- V. canadense Kalm. Sour-top or Velvet-leaf Blueberry. Cool moist woods; rare. May. Essex, (Jones).
- V. corymbosum L. High or Swamp Blueberry. Swamps and low thickets; common. May, June.

- V. macrocarpon Ait. Large or American Cranberry. Peat bogs; occasional. July, Aug.
- V. Oxycoccos L. Small Cranberry. Cold peat bogs; frequent. June, July. Var. INTERMEDIUM Gray. (Var. ovalifolium Michx.) Colchester, (Oakes); South Burlington, (Jones).
- V. pennsylvanicum Lam. Low Sweet or Early Sweet Blueberry. Dry hills, barrens, etc.; common. May.
- V. vacillans Kalm. Late Low Blueberry. Dry woodlands; common. May, June.

PRIMULACEAE. PRIMROSE FAMILY

ANAGALLIS (Tourn.) L. PIMPERNEL

A. ARVENSIS L. Common Pimpernel, Poor Man's Weatherglass. Adventive in garden, Burlington, (N. F. F.). June-Oct.

LYSIMACHIA (Tourn.) L. LOOSESTRIFE

- L. NUMMULARIA L. Moneywort, Yellow Myrtle. A frequent escape to damp grounds where it is thoroughly established. June, July.
- X L. producta (Gray) Fern. Mouth of Winooski River, Burlington, (Eggleston and Jones). July, Aug.
- L. quadrifolia L. Whorled Loosestrife. Dry thickets; common. June, July.
- L. terrestris (L.) BSP. Bulb-bearing Loosestrife. Low grounds; common. July, Aug.
- L. thyrsiflora L. Tufted Loosestrife. Bogs and swamps; frequent.

 June.

PRIMULA L. PRIMROSE

P. OFFICINALIS Jacq. English Cowslip. Roadside escape from garden, Burlington, (Jones). May.

STEIRONEMA Raf.

- S. ciliatum (L.) Raf. Fringed Loosestrife. Moist thickets; common. July, Aug.
- S. lanceolatum (Walt.) Gray. Lance-leaved Loosestrife. Low ground along Lake Champlain; rare. July, Aug. Colchester, (Grout and Tracy).

TRIENTALIS (Rupp.) L. CHICKWEED WINTERGREEN

T. americana (Pers.) Pursh. (T. borealis Raf.) Star Flower. Moist woods and thickets: common. May.

OLEACEAE. OLIVE FAMILY

FRAXINUS (Tourn.) L. ASH

- F. americana L. White Ash. Rich woods; common. May.
- F. nigra Marsh. Black Ash. Swamps and woods; frequent. May.
- F. pennsylvanica Marsh. Red Ash. Low grounds; common. May.
- F. pennsylvanica Marsh., var. lanceolata (Borkh.) Sarg. Green Ash. Shore of Lake Champlain, Burlington, (Jones). May.

SYRINGA L. LILAC

S. VULGARIS L. Common Lilac. Persistent about old building sites.

May, June.

GENTIANACEAE. GENTIAN FAMILY

BARTONIA Muhl.

B. virginica (L.) BSP. Yellow Bartonia. Boggy places; rare. July, Aug. South Burlington, (Jones, Ross); Colchester, (N. F. F.).

GENTIANA (Tourn.) L. GENTIAN

- G. Andrewsii Griseb. Closed Gentian. Moist grounds; frequent. Aug., Sept.
- G. quinquefolia L. Stiff Gentian, Ague-weed. Shore of Lake Champlain, South Burlington, (Torrey, Ross). Sept., Oct.

MENYANTHES (Tourn.) L. BUCKBEAN

M. trifoliata L. Bogs and shallow water; occasional. June.

APOCYNACEAE. DOGBANE FAMILY

APOCYNUM (Tourn.) L. DOGBANE. INDIAN HEMP

- A. androsaemifolium L. Spreading Dogbane. Thickets and fields; common. July-Sept.
- A. cannabinum L. Indian Hemp. Moist banks; frequent. July, Aug.

VINCA L. PERIWINKLE

V. MINOR L. Common Periwinkle, Blue Myrtle. Frequent in old gardens and established in rich woods near dwelling, Shelburne, (N. F. F.). Apr., May.

ASCLEPIADACEAE. MILKWEED FAMILY

ASCLEPIAS (Tourn.) L. MILKWEED. SILKWEED

- A. amplexicaulis Sm. Blunt-leaved Milkweed. Sandy soil; frequent. July.
- A. incarnata L. Swamp Milkweed. Marshes; common. July, Aug.
- A. phytolaccoides Pursh. Poke Milkweed. Moist copses; rare. July, Aug. Burlington, (Torrey, N. F. F.); Colchester, (N. F. F.).
- A. quadrifolia Jacq. Four-leaved Milkweed. Rocky woods; occasional. June, July.
- A. syriaca L. Common Milkweed or Silkweed. Rich grounds; common. July.

CONVOLVULACEAE. CONVOLVULUS FAMILY

CONVOLVULUS (Tourn.) L. BINDWEED

- C. ARVENSIS L. Field Bindweed. Railroad yard, Burlington, (Jones). July.
- C. JAPONICUS Thunb. California Rose. Occasional in old gardens and an escape by roadside, Essex, (Jones). July.
- C. sepium L. Wild Morning Glory, Hedge Bindweed. Moist thickets, sandy fields, etc.; common. July, Aug.
- C. spithamaeus L. Upright Bindweed. Dry sandy plains; common. June-Aug.

CUSCUTA (Tourn.) L. DODDER. LOVE VINE

C. Gronovii Willd. Moist thickets; frequent. July, Aug.

IPOMOEA L. MORNING GLORY

- I. COCCINEA L. Small Red Morning Glory. Adventive in garden, Burlington, (Mrs. Zottman). June-Aug.
- I. PURPUREA (L.) Roth. Common Morning Glory. Frequently spontaneous in gardens and waste places. June-Sept.

POLEMONIACEAE. POLEMONIUM FAMILY

GILIA R. & P.

G. LINEARIS (Nutt.) Gray. First seen in 1902, now well established in one station, Burlington, (N. F. F.). June, July.

PHLOX L.

- P. PANICULATA L. Garden Phlox. Occasionally persistent about old dwellings. Aug., Sept.
- P. Subulata L. Ground or Moss Pink. Rocky hill, Colchester, (Griffin). May, June.

HYDROPHYLLACEAE. WATERLEAF FAMILY

HYDROPHYLLUM (Tourn.) L. WATERLEAF

H. virginianum L. Rich woods; common. June.

BORAGINACEAE. BORAGE FAMILY

CYNOGLOSSUM (Tourn.) L. HOUND'S TONGUE

- C. boreale Fern. Wild Comfrey. Rich woods; rare. June. Burlington, (Torrey, Ross).
- C. OFFICINALE L. Common Hound's Tongue. Pastures; frequent. June, July.

ECHIUM (Tourn.) L. VIPER'S BUGLOSS

E. VULGARE L. Blue-weed, Blue Devil. Becoming frequent along railroads and occasional as a weed elsewhere. June-Oct.

LAPPULA (Rivin.) Moench. STICKSEED

- L. ECHINATA Gilib. European Stickseed, Burseed. Waste and cultivated grounds; occasional. Aug.
- L. virginiana (L.) Greene. Beggar's Lice. Borders of woods; occasional. Aug.

LITHOSPERMUM (Tourn.) L. GROMWELL, PUCCOON

L. latifolium Michx. Open ground and borders of woods; rare. June, July. Colchester, (Griffin).

I. officinale L. Common Gromwell. Dry pastures; occasional. June-Aug.

MERTENSIA Roth. LUNGWORT

M. VIRGINICA (L.) Link. Virginian Cowslip, Bluebells. A garden escape, Burlington, (Gifford). May. "Since this is a native from the Mississippi Valley reaching eastward to Ontario and New York, it may be expected to persist where established in the Champlain Valley."

MYOSOTIS (Rupp.) L. SCORPION-GRASS. FORGET-ME-NOT

M. SCORPIOIDES L. True Forget-me-not. Occasional on banks of Winooski River. June, July.

SYMPHYTUM (Tourn.) L. COMFREY

S. OFFICINALE L. Common Comfrey. A garden escape in moist ground, Shelburne, (N. F. F.). June-Sept.

VERBENACEAE. VERVAIN FAMILY

VERBENA (Tourn.) L. VERVAIN

- V. hastata L. Blue Vervain. Damp ground, roadsides, etc.; common. July-Sept.
- V. urticaefolia L. White Vervain. Roadsides; common. Aug., Sept.

LABIATAE. MINT FAMILY

AGASTACHE Clayt. GIANT HYSSOP

A. scrophulariaefolia (Willd.) Ktze. Thickets; rare. Aug., Sept. Essex, (N. F. F.).

AJUGA L. BUGLE WEED

A. GENEVENSIS L. Adventive in lawn, Burlington, (Jones). June.

GALEOPSIS L. HEMP NETTLE

G. Tetrahit L. Common Hemp Nettle. Waste places; frequent. July-Sept.

HEDEOMA Pers. Mock Pennyroyal

- H. HISPIDA Pursh. Adventive in garden, Burlington, (Hollister); Colchester, (Griffin). June, July.
- H. pulegioides (L.) Pers. American Pennyroyal. Dry fields; frequent. July, Aug.

LAMIUM (Tourn.) L. DEAD NETTLE

- L. AMPLEXICAULE L. Henbit. Cultivated ground; rare. June-Aug. South Burlington, (N. F. F.).
- L. MACULATUM L. Spotted Dead Nettle, Flowering Sage. Roadside escape from garden, Essex, (N. F. F.). May-Oct.

LEONURUS L. MOTHERWORT

L. CARDIACA L. Common Motherwort. Waste places; common. July, Aug.

LYCOPUS (Tourn.) L. WATER HOREHOUND

- L. americanus Muhl. Cut-leaved Bugle Weed. Moist soil; common. Aug., Sept.
- L. uniflorus Michx. Bugle Weed. Moist places; common. Aug., Sept.

MENTHA (Tourn.) L. MINT

- M. arvensis L., var. canadensis (L.) Briq. Wild Mint. Wet places; common. July, Aug.
- M. GENTILIS L. Spotted or Whorled Mint. Occasional in waste places and about old dwellings. July-Sept.
- M. PIPERITA L. Peppermint. Along brooks; rare. Aug., Sept. Essex and Shelburne, (N. F. F.); South Burlington, (Mrs. Walker).
- M. SPICATA L. Spearmint. Wet ground, Burlington, (N. F. F.). Aug., Sept.

MONARDA L. HORSE MINT

- M. MOLLIS L. Wild Bergamot. Dry grounds; rare. July, Aug. Burlington, (Jones); Shelburne, (Mrs. Gebhardt).
- M. PUNCTATA L. Horse Mint. Sandy ground; rare. July-Sept. Colchester and Burlington, (N. F. F.); Essex, (Jones and Howe).

NEPETA (Rivin.) L. CAT MINT

N. Cataria L. Catnip. Dry pastures; common. July-Sept.

N. HEDERACEA (L.) Trev. Ground Ivy, Gill-over-the-ground. Moist shady places; frequent. Apr., May.

PHYSOSTEGIA Benth. FALSE DRAGON HEAD

P. virginiana (L.) Benth. Lion's Heart. Wet ground; rare. July, Aug. Colchester, (Griffin).

PRUNELLA L. SELF-HEAL

P. vulgaris L. Heal-all, Carpenter-weed. Woods and fields; common. June-Sept. A white-flowered form is occasional.

PYCNANTHEMUM Michx. MOUNTAIN MINT. BASIL

- P. flexuosum (Walt.) BSP. Narrow-leaved Mountain Mint. Dry ground; rare. Aug. Williston, (N. F. F.); Colchester, (Ross).
- P. verticillatum (Michx.) Pers. Colchester, (Grout). Aug.
- P. virginianum (L.) Durand & Jackson. Dry woods, South Burlington, (N. F. F.). Aug.

SATUREJA (Tourn.) L. SAVORY. CALAMINT

- S. Acinos (L.) Scheele. Basil Thyme. Rocky headland of Lake Champlain, Burlington, (N. F. F.). June, July.
- S. HORTENSIS L. Summer Savory. A garden escape, Burlington, (Mrs. Walker). Aug., Sept.
- S. vulgaris (L.) Fritsch. Basil. Rocky woods and thickets; frequent. July-Sept.

SCUTELLARIA (Rivin.) L. SKULLCAP

- S. galericulata L. Marsh Skullcap. Wet shady places; frequent. July, Aug.
- S. lateriflora L. Mad-dog Skullcap. Wet shady places; common. July-Sept.
- S. parvula Michx. Small Skullcap. Rocky headlands of Lake Champlain; rare. June. Burlington and South Burlington, (Jones).

STACHYS (Tourn.) L. HEDGE NETTLE

S. tenuifolia Willd., var. aspera (Michx.) Fern. Rough Hedge Nettle, Woundwort. Moist grounds; occasional. July, Aug.

TEUCRIUM (Tourn.) L. GERMANDER

T. canadense L. American Germander, Wood Sage. Shores of Lake Champlain; frequent. Aug.

TRICHOSTEMA L. BLUE CURLS

T. dichotomum L. Bastard Pennyroyal. Sandy fields; local. Aug., Sept. Burlington, (N. F. F.); Colchester, (N. F. F., Griffin).

SOLANACEAE. NIGHTSHADE FAMILY

DATURA L. JAMESTOWN OF JIMSON WEED. THORN APPLE

- D. STRAMONIUM L. Stramonium. Waste places; occasional. July-Sept.
- D. TATULA L. Purple Thorn Apple. Occasionally adventive. July-Sept.

LYCIUM L. MATRIMONY VINE

L. HALIMIFOLIUM Mill. Common Matrimony Vine. An escape near waste ground, Burlington, (N. F. F.). May-Sept.

NICANDRA Adans. APPLE OF PERU

N. PHYSALODES (L.) Pers. Waste grounds; rare. July, Aug. Burlington, (Ross, N. F. F.).

PHYSALIS L. GROUND CHERRY

- P. HETEROPHYLLA Nees. One form occurs. Var. AMBIGUA (Gray) Rydb. Clammy Ground Cherry. Local. July-Sept. Burlington, (N. F. F., Jones).
- P. IXOCARPA Brot. Tomatillo. Adventive by roadside, Colchester, (Mrs. Walker). Aug., Sept.
- P. PERUVIANA L. Husk Tomato. Adventive on bank of Winooski River, Burlington, (Jones). Aug.
- P. PRUINOSA L. Strawberry Tomato. Adventive by roadside, Colchester, (Mrs. Walker). Aug., Sept.

SOLANUM (Tourn.) L. NIGHTSHADE

S. Dulcamara L. Bittersweet, Climbing Nightshade. Moist thickets; frequent. June-Aug.

- S. nigrum L. Common Nightshade. Moist shaded grounds; frequent. July-Sept.
- S. ROSTRATUM Dunal. Adventive in garden, Burlington, (Helyer). Aug.

SCROPHULARIACEAE. FIGWORT FAMILY

CHELONE (Tourn.) L. TURTLEHEAD. SNAKEHEAD

C. glabra L. Balmony. Wet places; frequent. Aug., Sept.

GERARDIA (Plum.) L. GERARDIA

- G. paupercula (Gray) Britton. Small-flowered Gerardia. Sandy shore of Lake Champlain, Colchester, (N. F. F., Griffin). Aug., Sept.
- G. tenuifolia Vahl. Slender Gerardia. Moist ground; local. Aug., Sept. Colchester, (Perkins, N. F. F., Ross).

GRATIOLA (Rupp.) L. HEDGE HYSSOP

- G. aurea Muhl. Golden Hedge Hyssop. Wet sandy shore of Lake Champlain, Colchester, (Perkins, N. F. F., Griffin). July, Aug.
- G. virginiana L. Clammy Hedge Hyssop. Wet shores and ditches; occasional. June-Sept.

ILYSANTHES Raf. FALSE PIMPERNEL

I. dubia (L.) Barnh. Short-stalked False Pimpernel. Wet shores; frequent. Aug., Sept.

LINARIA (Tourn.) Hill. TOADFLAX

L. VULGARIS Hill. Ramsted, Butter-and-Eggs. Fields and roadsides; common. July-Oct.

MELAMPYRUM (Tourn.) L. COW WHEAT

M. lineare Lam. Dry woodlands; frequent. July, Aug.

MIMULUS L. MONKEY FLOWER

M. ringens L. Wet places; common. July, Aug.

PEDICULARIS (Tourn.) L. LOUSEWORT

P. canadensis L. Common Lousewort, Wood Betony. Thickets; frequent. May, June.

PENTSTEMON (Mitch.) Ait. BEARD-TONGUE

P. LAEVIGATUS Ait., var. DIGITALIS (Sweet) Gray. Moist or rich soil; rare. June, July. Colchester, (Griffin); South Burlington, (Mrs. Walker).

SCROPHULARIA (Tourn.) L. FIGWORT

- S. leporella Bick. Meadows and fence rows; occasional. June, July.
- S. marilandica L. Damp grounds; occasional. July, Aug.

VERBASCUM (Tourn.) L. MULLEIN

- V. BLATTARIA L. Moth Mullein. Roadsides; rare. June, July. Burlington and Essex, (N. F. F.).
- V, THAPSUS L. Common Mullein. Fields, etc.; common. July-Sept.

VERONICA (Tourn.) L. SPEEDWELL

- V. americana Schwein. American Brooklime. Wet grounds; frequent. June, July.
- V. ARVENSIS L. Corn Speedwell. Rocky woods; frequent. May, June.
- V. officinalis L. Common Speedwell. Dry hills and open woods; occasional. May, June.
- V. PEREGRINA L. Neckweed, Purslane Speedwell. Gardens and waste places; occasional. May, June.
- V. scutellata L. Marsh Speedwell. Swamps and wet places; common. June-Aug.
- V. serpyllifolia L. Thyme-leaved Speedwell. Damp grassy grounds; common. May-July.

LENTIBULARIACEAE. BLADDERWORT FAMILY

UTRICULARIA L. BLADDERWORT

- U. intermedia Hayne. Edge of swamp, Colchester, (N. F. F.). June, July.
- U. vulgaris L., var. americana Gray. Greater Bladderwort. Ponds and slow streams; frequent. July, Aug.

OROBANCHACEAE. BROOM=RAPE FAMILY

EPIFAGUS Nutt. BEECH-DROPS. CANCER-ROOT

E. virginiana (L.) Bart. Under beech trees; common. Sept.

OROBANCHE (Tourn.) L. BROOM-RAPE

 uniflora L. One-flowered Cancer-root. Dry woodlands; rare. May-July. Burlington, (Jones, Mrs. Walker).

PHRYMACEAE. LOPSEED FAMILY

PHRYMA L. LOPSEED

P. Leptostachya L. Moist open woods; occasional. July, Aug.

PLANTAGINACEAE. PLANTAIN FAMILY

PLANTAGO (Tourn.) L. PLANTAIN. RIBWORT

- P. ARISTATA Michx. Large-bracted Plantain. Adventive at Experiment Farm, Burlington, (Jones); dry sand, Colchester, (N. F. F.). July, Aug.
- P. LANCEOLATA L. Rib or Ripple Grass, English Plantain. Grass land; common. June-Aug.
- P. major L. Common Plantain. Waysides; common. July-Sept.
- P. Rugelii Done. With the preceding species and even more common. July-Sept.

RUBIACEAE. MADDER FAMILY

CEPHALANTHUS L. BUTTONBUSH

C. occidentalis L. Swamps; frequent. July, Aug.

GALIUM L. BEDSTRAW. CLEAVERS

- 6. Aparine L. Cleavers, Goose Grass. Rich shaded grounds; occasional. May, June. Burlington and South Burlington, (N. F. F.); Colchester, (Griffin).
- G. asprellum Michx. Rough Bedstraw. Alluvial grounds; common. July,

- G. BOREALE L. Northern Bedstraw. Edge of woods, Burlington, (N. F. F.). July.
- G. circaezans Michx. Wild Liquorice. Rich woods; common. July.
- G. Claytoni Michx. Swamps and bogs; occasional. July-Sept. Burlington and Colchester, (N. F. F.).
- G. lanceolatum Torr. Wild Liquorice. Dry woods; frequent. July.
- G. Mollugo L. Wild Madder. Adventive in field, Shelburne, (N. F. F.).
 June, July.
- G. palustre L. Marsh Bedstraw. Wet meadows and banks; common. June, July.
- G. trifidum L. Small Bedstraw. Cold bogs and marshy borders of ponds; frequent. July, Aug.
- G. triflorum Michx. Sweet-scented Bedstraw. Rich woodlands; common. July, Aug.

HOUSTONIA L.

- H. caerulea L. Bluets, Innocence. Moist grassy places; common. Apr., May. The white flowered form occurs and occasional blossoms may be found all summer.
- H. longifolia Gaertn. Rocky headlands of Lake Champlain; rare. June.

MITCHELLA L. PARTRIDGE BERRY

M. repens L. Cool woods; common. June. A form occurs the leaves of which are variegated with white; Burlington, (N. F. F.).

CAPRIFOLIACEAE. HONEYSUCKLE FAMILY

DIERVILLA (Tourn.) Mill. BUSH HONEYSUCKLE

D. Lonicera Mill. Rocky woodlands; common. June, July.

LINNAEA (Gronov.) L. TWIN-FLOWER

L. borealis L., var. americana (Forbes) Rehder. Moist mossy woods and cool bogs; rare. June, July. Shelburne, (Jones); South Burlington, (N. F. F.).

LONICERA L. HONEYSUCKLE

L. canadensis Marsh. American Fly Honeysuckle. Moist woods; frequent. Apr., May.

- L. dioica L. Glaucous Honeysuckle. Dry rocky and sandy woods; frequent. May.
- L. TATARICA L. Tartarian Honeysuckle. Common in cultivation and a frequent escape in thickets about Burlington. May.

SAMBUCUS (Tourn.) L. ELDER

- S. canadensis L. Common Elder, Elderberry. Thickets; common. July.
- S. racemosa L. Red-berried Elder. Rocky woodlands; common. May.

SYMPHORICARPOS (Dill.) Ludwig. SNOWBERRY

- S. racemosus Michx. Snowberry. Cliffs and rocky headlands; frequent. June.
- S. RACEMOSUS Michx., var. LAEVIGATUS Fern. Garden Snowberry. Common in cultivation and occasionally persistent about old building sites. June, July. Shelburne, (N. F. F.).

TRIOSTEUM L. FEVERWORT. HORSE GENTIAN

T. perfoliatum L. Tinker's Weed, Wild Coffee. Edge of swamp, Colchester, (Griffin). May, June.

VIBURNUM (Tourn.) L. ARROW-WOOD, LAURESTINUS

- V. acerifolium L. Dockmackie, Maple-leaved Arrow-wood. Rocky or sandy woods; common. June.
- V. alnifolium Marsh. Hobble-bush, Witch Hobble, Moosewood. Cool moist woods: common. May.
- V. cassinoides L. Withe-rod, Wild Raisin. Swamps; frequent. June.
- V. dentatum L. Toothed Arrow-wood. Wet places; rare. June, July. Essex and Shelburne, (N. F. F.).
- V. Lentago L. Sheepberry, Nannyberry, Wild Raisin. Rich moist woods; occasional. May, June.
- V. Opulus L., var. americanum (Mill.) Ait. Cranberry Tree, High-bush Cranberry, Pimbina. Low grounds; occasional. May, June.
- V. pubescens (Ait.) Pursh. Downy Arrow-wood. Cliffs and rocky hills; frequent. June.

VALERIANACEAE. VALERIAN FAMILY

VALERIANA (Tourn.) L. VALERIAN

V. OFFICINALIS L. Garden Heliotrope. An occasional roadside escape.

June. Burlington and Shelburne, (N. F. F.).

CUCURBITACEAE. GOURD FAMILY

ECHINOCYSTIS T. & G. WILD BALSAM-APPLE

E. lobata (Michx.) T. & G. Wild Cucumber. Alluvial banks and waste places; frequent. July-Sept.

SICYOS L. ONE-SEEDED BUR CUCUMBER

S. angulatus L. Star Cucumber. River banks and waste places; frequent. July-Sept.

CAMPANULACEAE. BLUEBELL FAMILY

CAMPANULA (Tourn.) L. BELLFLOWER

- C. aparinoides Pursh. Marsh Bellflower. Low sedgy thickets; occasional. July, Aug.
- C. RAPUNCULOIDES L. European Bellflower. An occasional roadside escape. July, Aug.
- C. rotundifolia L. Harebell, Bluebell. Rocky banks; common. June-Oct.
- C. TRACHELIUM L. Nettle-leaved Bellflower, Throatwort. Adventive, Colchester, (Griffin). Aug.

SPECULARIA (Heist.) Fabric. VENUS'S LOOKING-GLASS

S. perfoliata (L.) DC. Dry rocky woodlands; rare. June, July. South Burlington, (N. F. F.).

LOBELIACEAE. LOBELIA FAMILY

LOBELIA (Plum.) L.

- L. cardinalis L. Cardinal Flower. Low grounds; occasional. Aug.
- L. inflata L. Indian Tobacco. Fields and pastures; common. July-Sept.

- L. Kalmii L. Cool moist rocks; occasional. July, Aug.
- L. spicata Lam. Moist sandy soil; occasional. July, Aug.

COMPOSITAE. COMPOSITE FAMILY

ACHILLEA (Vaill.) L. YARROW

A. Millefolium L. Common Yarrow, Milfoil. Dry fields and roadsides; common. July-Oct. The pink-flowered form is occasional.

AMBROSIA (Tourn.) L. RAGWEED

- A. artemisiifolia L. Roman Wormwood, Hog-weed, Bitter-weed. A common weed everywhere. Aug., Sept.
- A. PSILOSTACHYA DC. Adventive in railroad yard, Burlington, (N. F. F.). Aug., Sept.
- A. trifida L. Great Ragweed. Moist alluvial soil; local. Williston, (Bates); Burlington, (N. F. F.). Rare as a weed elsewhere. Aug. Var. INTEGRIFOLIA (Muhl.) T. & G. Burlington, (N. F. F.)

ANAPHALIS DC. EVERLASTING

- A. margaritacea (L.) B. & H. Pearly Everlasting. Dry hills and woods; common. July-Sept.
- ANTENNARIA Gaertn. Everlasting. Ladies' Tobacco. Pussy's Toes
- A. canadensis Greene. Dry and moist ground; frequent. May, June.
- A. fallax Greene. In partial shade; frequent. May.
- A. neglecta Greene. Old fields; frequent. May.
- A. neodioica Greene. Old pastures and fields; frequent. May, June.
- A. Parlinii Fern. Dry banks and thickets; rare. May, June. Colchester, (Griffin).
- A. petaloidea Fern. Old pastures; rare. May. Shelburne, (Brainerd); Burlington, (Hazen, N. F. F.).
- A. plantaginifolia (L.) Richards. Plantain-leaved Everlasting. Dry soil; common. May, June.

ANTHEMIS (Mich.) L. CHAMOMILE

A. COTULA L. Mayweed, Dog Fennel. Waste places and roadsides; common. June-Oct.

ARCTIUM L. BURDOCK

- A. LAPPA L. Great Burdock. Roadsides; local. July-Sept. Essex, (Jones); Colchester, (N. F. F.).
- A. MINUS (Hill) Bernh. Common Burdock. Waste places; common. July-Oct.

ARTEMISIA (Tourn.) L. WORMWOOD

- A. Absinthium L. Wormwood. Roadside, Colchester, (N. F. F.).

 Aug., Sept.
- A. BIENNIS Willd. Abundant around railroad and lumber yards, Burlington. Aug., Sept.
- A. caudata Michx. Tall Wormwood. Abundant on sandy bluff of Lake Champlain, Burlington. Aug., Sept.
- A. VULGARIS L. Common Mugwort. Waste places and roadsides; common. July, Aug.

ASTER (Tourn.) L. ASTER

- A. acuminatus Michx. Cool rich woods; frequent. July, Aug.
- A. cordifolius L. Heart-leaved Aster. Woodlands; common. Aug., Sept.
- A. divaricatus L. White Wood Aster. Moist woodlands; frequent. Aug., Sept.
- A. ericoides L. One form occurs. Var. Pringlei Gray. Pringle's Aster. Slaty headlands of Lake Champlain; rare. Aug. Shelburne, (Pringle); Burlington.
- A. lateriflorus (L.) Britton. Thickets and fields; occasional. Aug., Sept.
- A. lateriflorus (L.) Britton, var. hirsuticaulis (Lindl.) Porter. Hairy-stemmed Aster. Borders of thickets; common. Aug., Sept.
- A. linariifolius L. Stiff Aster. Dry sandy soil; common. Aug., Sept.
- A. macrophyllus L. Large-leaved Aster. Moist woods; common. Aug., Sept. Presenting many different forms.
- A. novae-angliae L. New England Aster. Moist grounds; rare. Sept. Burlington, (Jones, N. F. F.). Var. ROSEUS (Desf.) DC. Burlington, (Miss Moore).
- A. paniculatus Lam. Tall White or Panicled Aster. Moist soil; common. Sept. Presenting many different forms.

- A. puniceus L. Purple-stemmed Aster. Low thickets and swamps; frequent. Sept.
- A. Tradescanti L. Michaelmas Daisy. Low grounds; common. Sept.
- A, umbellatus Mill. Flat-topped White Aster. Moist thickets; frequent. Aug., Sept.
- A. undulatus L. Wavy-leaved Aster. Dry copses; common. Aug., Sept. Variable.
- A. vimineus Lam. Small White Aster. Moist banks; occasional. Aug.
- A. vimineus Lam., var. saxatilis Fern. Ledges of Winooski River, Williston, (Pringle); Burlington, (N. F. F.). Aug.

BIDENS L. BUR MARIGOLD

- B. Beckii Torr. Water Marigold. Ponds and deep slow streams; rare. Aug. Shelburne, (Jones); Colchester, (N. F. F., Mrs. Walker).
- B. cernua L. Stick-tight. Wet places; common. Aug., Sept.
- B. connata Muhl. Swamp Beggar-ticks. Swamps; occasional. Aug., Sept.
- S. frondosa L. Common Beggar-ticks. Damp grounds; common. Aug., Sept.

CARDUUS (Tourn.) L. PLUMELESS THISTLE

C. CRISPUS L. Adventive, South Burlington, (Chittenden). Aug.

CENTAUREA L. STAR THISTLE

C. Cyanus L. Blue Bottle, Bachelor's Button, Corn Flower. Spontaneous in gardens and occasionally adventive elsewhere. June-Sept.

CHRYSANTHEMUM (Tourn.) L. OX-EYE DAISY

C. LEUCANTHEMUM L., var. PINNATIFIDUM Lecoq & Lamotte. Ox-eye or White Daisy, Marguerite, Whiteweed. A common and pernicious weed in fields and meadows. June-Oct.

CICHORIUM (Tourn.) L. SUCCORY OF CHICORY

C. Intybus L. Common Chicory, Blue Sailors. Roadsides and fields; common in Burlington, less so elsewhere. July, Aug.

CIRSIUM (Tourn.) Hill. COMMON OF PLUMED THISTLE

- C. ARVENSE (L.) Scop. Canada Thistle. Fields and waste places; common. June-Sept.
- C. LANCEOLATUM (L.) Hill. Common or Bull Thistle. Pastures and roadsides; common. July-Sept.

DYSSODIA Cav. FETID MARIGOLD

D. PAPPOSA (Vent.) Hitchc. Adventive on wool waste, bank of Winooski River, Burlington, (Brainerd). July-Oct.

ERECHTITES Raf. FIREWEED

E. hieracifolia (L.) Raf. Moist woods, especially in recent clearings; common. July-Sept.

ERIGERON L. FLEABANE

- E. annuus (L.) Pers. Daisy Fleabane, Sweet Scabious. Fields and waste places; common. June-Sept.
- E. cauadensis L. Horseweed, Butterweed. Waste places, etc.; common. July-Sept.
- E. hyssopifolius Michx. Wet calcareous rocks; occasional. June, July.
- E. philadelphicus L. Pink Fleabane. Moist grounds; common. May-July.
- E. pulchellus Michx. Robin's Plantain. Copses and moist banks; frequent. May, June.
- E. ramosus (Walt.) BSP. Daisy Fleabane. Fields; common. June-Sept.

EUPATORIUM (Tourn.) L. THOROUGHWORT

- E. perfoliatum L. Thoroughwort, Boneset. Low grounds; common. Aug., Sept.
- E. purpureum L., var. maculatum (L.) Darl. Joe-Pye Weed, Trumpet Weed. Moist thickets and low grounds; common. July, Aug.
- E. urticaefolium Reich. White Snakeroot. Moist rich woods; occasional. Aug., Sept.

GALINSOGA R. & P.

G. PARVIFLORA Cav. One form occurs. Var. HISPIDA DC. First collected in 1891, (Grout), now a common and pernicious garden weed. July-Oct.

GNAPHALIUM L. CUDWEED

- G. decurrens Ives. Everlasting. Clearings, etc.; frequent. Aug., Sept.
- G. polycephalum Michx. Common Everlasting. Old fields and woods; common. Aug., Sept.
- G. uliginosum L. Low Cudweed. Moist soil; common. July-Sept.

HELIANTHUS L. SUNFLOWER

- II. ANNUUS L. Common Sunflower. Occasionally spontaneous in waste places. Aug., Sept.
- H. decapetalus L. Thin-leaved Sunflower. Copses and low banks of streams; occasional. Aug., Sept.
- H. divaricatus L. Rough or Woodland Sunflower. Thickets and sandy woods; common. July-Sept.
- H. PETIOLARIS Nutt. Adventive in garden, Burlington, (Helyer). Aug., Sept.
- H. strumosus L. Pale-leaved Wood Sunflower. Dry woods and banks; frequent. Aug., Sept.
- H. Tuberosus L. Jerusalem Artichoke. Along fences and roadsides; frequent. Sept., Oct.

HELIOPSIS Pers. OX-EYE

H. SCABRA Dunal. Adventive in fields and by roadsides. Aug., Sept. Burlington, (Jones); Shelburne, (Mrs. Gebhardt).

HIERACIUM (Tourn.) L. HAWKWEED

- H. AURANTIACUM L. Orange Hawkweed, Devil's Paint-brush, Grim the Collier. Fields and pastures; a common and pernicious weed. June-Sept.
- H. canadense Michx. Canada Hawkweed. Borders of woods; frequent. Aug., Sept.
- H. FLORENTINUM All. King Devil. One plant on railroad embankment, Burlington, (N. F. F.); Colchester, (Griffin). June.
- H. paniculatum L. Panicled Hawkweed. Open woods; occasional. Aug.
- H. PRAEALTUM Goch., var. DECIPIENS Koch. Interval land, South Burlington, (Mrs. Walker). June.

- H. scabrum Michx. Rough Hawkweed. Dry plains and pine woods; frequent. July, Aug.
- H. venosum L. Rattlesnake-weed, Poor Robin's Plantain. Dry plains and pine woods; frequent. June, July.

INULA L. ELECAMPANE

I, HELENIUM L. Elecampane. Roadsides and damp pastures; frequent. Aug.

•LACTUCA (Tourn.) L. LETTUCE

- L. canadensis L. Wild Lettuce, Horseweed. Rich damp soil, borders of fields and thickets; frequent. July, Aug. Var. Montana Britton. South Burlington, (N. F. F.). July, Aug.
- L. SCARIOLA L., var. INTEGRATA Gren. & Godr. Prickly Lettuce. Abundant along Rutland R. R. in some localities in Burlington and Colchester, and occasional as a weed elsewhere. July-Sept.
- L. spicata (Lam.) Hitchc. Tall Blue Lettuce. Low grounds; frequent. July, Aug.

MATRICARIA (Tourn.) L. WILD CHAMOMILE

M. SUAVEOLENS (Pursh) Buch. Pine-apple-weed. About waste ground and railroad yards, Burlington, (N. F. F.). June, July.

PRENANTHES (Vaill.) L. RATTLESNAKE-ROOT

- P. alba L. White Lettuce, Rattlesnake-root. Borders of rich woods; common. Aug., Sept.
- P. altissima L. Tall White Lettuce. Rich moist woods; frequent. Aug., Sept.
- P. trifoliolata (Cass.) Fern. Gall-of-the-earth. Thickets and woods; rare. Aug., Sept. Burlington, (Grout); Essex, (N. F. F.); Colchester, (Griffin).

RUDBECKIA L. CONE-FLOWER

- R. HIRTA L. Yellow Daisy, Black-eyed Susan, Nigger-head. Meadows; common. June-Sept.
- R. laciniata L. Tall Cone-flower, Thimble-weed. Low thickets; rare. Aug., Sept. High Bridge, (Jones); Burlington, (N. F. F.).

SENECIO (Tourn.) L. GROUNDSEL. RAGWORT. SQUAW-WEED

- S. aureus L. Golden Ragwort. Moist rocks, High Bridge, (Ross).
 June, July.
- S. Balsamitae Muhl. Balsam Groundsel. Rocky banks of Winooski River; frequent. May-July.
- S. Balsamitae Muhl., var. praelongus Greenm. Rocky bank of Winooski River near Twin Bridges, (Jones); Essex, (Ross). June.
- S. obovatus Muhl. Round-leaved Squaw-weed. Rocky bank of Winooski River, Essex, (Ross). June.
- S. Robbinsii Oakes. Robbin's Squaw-weed. Wet meadows and swamps; rare. June, July. South Burlington, (Ross); Shelburne, (N. F. F.).
- S. VULGARIS L. Common Groundsel. A weed in gardens and waste places; occasional. July-Sept.

SILPHIUM L. ROSIN-WEED

S. PERFOLIATUM L. Cup Plant. Adventive by roadside, Burlington, (Jones). Aug., Sept.

SOLIDAGO L. GOLDEN-ROD

- S. arguta Ait. Cut-leaved Golden-rod. Burlington, (Grout). Aug.
- S. bicolor L. White Golden-rod. Dry copses; common. Aug., Sept.
- S. caesia L. Blue-stemmed Golden-rod. Rich woodlands; common. Aug., Sept.
- S. canadensis L. Canada Golden-rod. Borders of thickets and fields; common. Aug., Sept.
- S. graminifolia (L.) Salisb. Bushy Golden-rod. Moist soil; common. Aug., Sept.
- S. hispida Muhl. Hairy Golden-rod. Dry shady cliffs and sterile soil; occasional. Aug., Sept.
- S. juncea Ait. Early Golden-rod. Copses and banks; common. July, Aug.
- S. latifolia L. Zig-zag Golden-rod. Moist shaded banks; frequent. Aug., Sept.
- S. nemoralis Ait. Gray or Field Golden-rod. Dry open soil; common.

 Aug., Sept. A peculiar low form is occasional with the type.

- S. puberula Nutt. Downy Golden-rod. Sandy soil; common. Aug., Sept.
- S. racemosa Greene. Rocky banks of Winooski River; occasional. Aug. Winooski Gorge is the station of Pursh's type.
- S. rugosa Mill. Tall Hairy Golden-rod. Borders of fields and copses; common. Aug., Sept.
- S. serotina Ait. Late Golden-rod. Thickets, in rich soil; frequent. Aug., Sept.
- S. squarrosa Muhl. Stout Ragged Golden-rod. Dry woods; frequent. Aug., Sept.
- S. uliginosa Nutt. Bog Golden-rod. Peat bogs; rare. Sept. South Burlington, (Jones).

SONCHUS (Tourn.) L. Sow Thistle

- S. ARVENSIS L. Field Sow Thistle. Shores and waste places; occasional. July-Sept.
- S. ASPER (L.) Hill. Spiny-leaved Sow Thistle. Waste places; occasional. July-Oct.
- S. OLERACEUS L. Common Sow Thistle. Waste places; frequent. July-Oct.

TANACETUM (Tourn.) L. TANSY

T. VULGARE L. Common Tansy. Roadsides, etc.; frequent. Aug., Sept. Var. CRISPUM DC. South Burlington, (Ross); Burlington, (N. F. F.).

TARAXACUM (Haller) Ludwig. Dandelion

- T. ERYTHROSPERMUM Andrz. Red-seeded Dandelion. Shallow soil about ledges; frequent. Apr., May.
- T. OFFICINALE Weber. Common Dandelion. Common everywhere. Apr.-Sept.

TRAGOPOGON (Tourn.) L. GOAT'S BEARD

T. PRATENSIS L. Roadsides and meadows; occasional. June-Sept.

TUSSILAGO (Tourn.) L. COLTSFOOT

T. FARFARA L. Moist clay banks; occasional. Apr., May.

XANTHIUM (Tourn.) L. COCKLEBUR. CLOTBUR

- X. canadense Mill. Lake shores and river banks; frequent. Aug., Sept.
- X. commune Britton. Shores of Lake Champlain; common. Aug., Sept.
- X. echinatum Murr. Lake shores and river banks; common. Aug., Sept.

SUPPLEMENTARY LIST

This is a list of plants which should be sought in the region covered by this Flora, and comprises those plants which are known to have occurred here, but of which no specimen was taken or preserved, those which occur near the limits of this region and may naturally be expected here, and those specimens of which are in the Torrey collection, U. V. M. herbarium, but without locality.

Others have been taken from Perkins' Flora, Thompson's Vermont and an old list of Burlington plants published in the Burlington Free Press about 1845, in the hope that search may reveal some of them.

- Osmunda cinnamomea L., var. frondosa Gray. Swamps and low copses.

 May. Charlotte, (Pringle).
- Isoetes sp. In water and wet places. Some of these ought to be found here.
- Sparganium simplex Huds. In water. July. Charlotte, (Pringle).
- Potamogeton alpinus Balbis. Ponds and streams. July, Aug. Hinesburgh Pond, (Jones).
- Potamogeton obtusifolius Mertens & Koch. Ponds and streams. July, Aug. Ferrisburgh, (Grout and Eggleston).
- Potamogeton angustifolius Berchtold & Presl. Lake Champlain, Milton, (Grout).
- Ammophila arenaria (L.) Link. Sandy beaches. Aug., Sept. Reported from Colchester Point, (Jones).
- Deschampsia caespitosa (L.) Beauv. Reported from rocky shore of Winooski River, south side, between Essex Junction and Burlington, (Jones). He says the species was undoubted, but no herbarium specimen can be found. June, July.
- Holcus lanatus L. Fields and meadows. June, July. Charlotte, (Pringle).
- Milium effusum L. Cold damp woods. June. Torrey collection.
- Muhlenbergia sylvatica Torr. Moist rocky woods and wooded banks.

 Aug., Sept. Torrey collection. Ferrisburgh, (Pringle).

- Carex aquatilis Wahl. Swamps and lake margins. June-Aug. Torrey collection.
- Carex Bicknellii Britton. Dry rocky soil. June, July. Torrey collection.
- Carex flava L., var. rectirostris Gaudin. Wet margins of springs and low borders of ponds. July, Aug. Torrey collection.
- Carex gynocrates Wormsk. Swamps and bogs. June-Aug. Burlington, (Torrey); Perkins' Flora.
- Carex polygama Schkuhr. Bogs and wet shores. May-July. Torrey collection.
- Carex retrorsa Schwein., var. Hartii (Dewey) Gray. Wet places. July, Aug. Torrey collection.
- Carex triceps Michx., var. hirsuta (Willd.) Bailey. Moist fields. June, July. Charlotte, (Pringle).
- Carex trichocarpa Muhl. Marshes. June, July. Burlington; Perkins' Flora.
- Carex retroflexa Muhl. Dry open woods. May, June. Burlington, (Torrey); Perkins' Flora.
- Carex vaginata Tausch. Bogs and mossy woods. June, July. Torrey collection.
- Carex virescens Muhl. Moist pastures. June, July. Charlotte, (Pringle).
- Scirpus planifolius Muhl. Dry open fields. May, June. Charlotte, (Pringle).
- Peltandra virginica (L.) Kunth. Bogs. July. Colchester, (Robbins); Thompson's Vermont.
- Juneus filiformis L. Wet shores and bogs. June, July. Shore of Lake Champlain, Ferrisburgh, (Robbins); Thompson's Vermont.
- Zigadenus chloranthus Richards. Calcareous soils. July. Ferrisburgh, (Brainerd).
- Hypoxis hirsuta (L.) Coville. Meadows and open woods. June, July. Free Press.
- Sisyrinchium gramineum Curtis. Wet meadows and damp woods. May, June. Burlington, (Macrae); Thompson's Vermont.
- Corallorrhiza odontorhiza Nutt. Woods. Aug., Sept. Torrey collection. Reported from Mud Pond, Williston, (Bryant).

- Corallorrhiza striata Lindl. Woods. May, June. Torrey collection.
- Habenaria flava (L.) Gray. Wet places. June, July. Burlington, (Macrae); Thompson's Vermont.
- Liparis liliifolia (L.) Richards. Woods. June, July. Found just south of Fort Ethan Allen, but specimen lost.
- Spiranthes lucida (Eaton) Ames. Moist banks. June, July. Torrey collection. Reported from near Winooski Gorge, (Ross).
- Carya glabra (Mill.) Spach. Dry woodlands. May. Torrey collection. Ferrisburgh, (Brainerd).
- Corylas americana Walt. Reported from sand plains, South Burlington, (Howe).
- Quercus Prinns L. Rocky banks and hillsides. May. Charlotte, (Pringle).
- Ulmus racemosa Thomas. Rocky woodlands. May. Occurs as a street tree in Burlington.
- Polygonum amphibium L. Margins of ponds and slow streams. Aug., Sept. Richmond, (Pringle).
- Polygonum Douglasii Greene. Rocky or sterile soil. Aug., Sept. Charlotte, (Pringle).
- Polygonum hydropiperoides Michx. Wet places and in shallow water. Aug., Sept. Charlotte, (Pringle).
- Polygonum tenue Michx. Dry soil. Aug., Sept. Torrey collection. This occurs at Keeseville on opposite side of Lake Champlain.
- Amaranthus paniculatus L. Weed in garden, Shelburne, (Pringle); Flora of Vermont.
- ('eratophyllum demersum L. Slow streams and ponds. June, July. Reported from Ferrisburgh. Mouth of Winooski River, (Robbins); Thompson's Vermont.
- Anemonella thalietroides (L.) Spach. Woods. May. Reported from Burlington, (Benedict).
- Cimicifuga racemosa (L.) Nutt. Rich woods. July. Torrey collection.
- Ranunculus fascicularis Muhl. Dry or moist hills. Apr., May. Reported from Burlington, (Benedict, Bigelow).
- Menispermum canadense L. Banks of streams. June, July. Charlotte, (Pringle). Reported from Ethan Allen Park, Burlington, (Ross). Burlington, (Carey); Thompson's Vermont.

Arabis canadensis L. Rocky woods and ravines. June-Aug. Torrey collection. Winooski River, Colchester, (Torrey); Perkins' Flora.

Arabis lyrata L. Rocks or sandy shores. May-July. Torrey collection.

Cardamine bulbosa (Schreb.) BSP. Wet meadows. May. Torrey collection.

Radicula Nasturtium-aquaticum (L.) Britten & Rendle. Brooks, ditches, etc. June-Aug. Charlotte, (Pringle).

Crataegus anomala Sarg. Fl. May; fr. Oct.

Crataegus Brainerdi Sarg., var. asperifolia (Sarg.) Eggl.

Crataegus Brainerdi Sarg., var. Egglestoni (Sarg.) Robins.

Crataegus Brainerdi Sarg., var. scabrida (Sarg.) Eggl.

Crataegus champlainensis Sarg. Fl. May; fr. Sept., Oct.

Crataegus Holmesiana Ashe. Fl. May; fr. Aug., Sept.

Crataegus macracantha Lodd, var. neotluvialis (Ashe) Eggl. Fl. May; fr. Sept.

Crataegus macracantha Lodd, var. succulenta (Schrad.) Eggl. Fl. May; fr. Sept.

Crataegus macrosperma Ashe, var. demissa (Sarg.) Eggl. Fl. May; fr. Sept.

Crataegus macrosperma Ashe, var. pastorum (Sarg.) Eggl. Fl. May; fr. Sept.

Crataegus macrosperma Ashe, var. pentandra (Sarg.) Eggl. Fl. May; fr. Sept.

Crataegus polita Sarg. Fl. May; fr. Sept.

Crataegus Pringlei Sarg. Fl. May; fr. Sept. Charlotte, (Pringle).

Crataegus Pringlei Sarg., var. exclusa (Sarg.) Eggl. Fl. May; fr. Sept.

Crataegus Pringlei Sarg., var. lobulata (Sarg.) Eggl. Fl. May; fr. Sept.

Crataegus pruinosa (Wendl.) Koch forma dissona (Sarg.) Eggl. Fl.
May; fr. Oct.

Crataegus punctata Jacq., var. canescens Britton. Fl. May; fr. Oct.

Crataegus punctata Jacq., var. rubra Ait. Fl. May, June; fr. Oct.

Crataegus roanensis Ashe. Fl. May; fr. Sept.

- Crataegus rotundifolia Moench, var. Faxoni (Sarg.) Eggl. Fl. May; fr. Aug., Sept.
- Desmodium paniculatum (L.) DC. Copses. July. Charlotte, (Pringle).
- Lespedeza frutescens (L.) Britton. Open rocky woods. Aug., Sept. Mt. Philo, Charlotte, (Pringle).
- Lespedeza hirta (L.) Hornem. Dry hills and plains. Aug., Sept. Colchester, (Robbins); Thompson's Vermont.
- Oxalis violacea L. Rocky places and open woods. May. Free Press.
- Euphorbia Peplus L. Waste places and cultivated grounds. Aug. Charlotte, (Pringle).
- Euphorbia Preslii Guss. Dry open soil. July, Aug. Burlington, (Tuckerman); Thompson's Vermont.
- Rhus canadensis Marsh. Dry rocky banks. May. Charlotte, (Pringle).
- **Hypericum majus** (Gray) Britton. Wet or dry open soil. July, Aug. Charlotte, (Pringle).
- Conioselinum chinense (L.) BSP. Swamps and cold cliffs. Aug., Sept. Burlington, (Macrae); Thompson's Vermont.
- Cornus florida L. Dry woods. May. Free Press.
- Lyonia ligustrina (L.) DC. Moist thickets. June, July. Torrey collection.
- Pyrola secunda L., var. obtusata Turcz. Reported from South Burlington, (Ross).
- Rhododendron canescens (Michx.) G. Don. Woods and gravelly shores.

 May, June. Reported from Burlington, (N. F. F.).
- Rhododendron viscosum (L.) Torr. Swamps. June, July. Free Press.
- Vaccinium corymbosum L., var. amoenum (Ait.) Gray. Swamps and low thickets. May, June. Charlotte, (Pringle).
- Gentiana crinita Froel. Low grounds. Sept. Torrey collection.
- Gentiana linearis Froel. Bogs and wet rocks. Aug. Torrey collection. Hinesburgh, (Horsford).
- Halenia deflexa (Sm.) Griseb. Damp and cool woods. Aug., Sept. Charlotte, (Horsford and Pringle).
- Asclepias incarnata L., var. pulchra (Ehrh.) Pers. Swamps. July, Aug. Torrey collection.

- Lithospermum arvense L. Sandy fields and roadsides. May-Aug. Charlotte, (Pringle).
- Lycopsis arvensis L. Dry or sandy fields and waste places. July. Torrey collection.
- **Pycnanthemum muticum** (Michx.) Pers. Reported from bank of Winooski River, Colchester, (Grout).
- Gerardia flava L. Open woods. July, Aug. Torrey collection.
- Gerardia pedicularia L. Dry copses. July, Aug. Torrey collection.
- Pentstemon hirsutus (L.) Willd. Dry or rocky grounds. June, July. Charlotte, (Pringle). Torrey collection.
- Veronica Anagallis-aquatica L. Brooks and ditches. June-Aug. Reported from Williston, (Bryant).
- Veronica virginica L. Rich soil. July, Aug. Colchester, (Oakes); Thompson's Vermont.
- Conopholis americana (L. f.) Wallr. Rock Pt., Burlington, (Macrae); Thompson's Vermont.
- Dianthera americana L. In water. July, Aug. Ferrisburgh, (Paddock).
- Littorella uniflora (L.) Asch. In water or on gravelly shores. Aug., Sept. "Ought to be found here on lake shore." Pringle.
- Dipsaeus sylvestris Huds. Roadsides. July, Aug. Charlotte, (Pringle).
- Lobelia Dortmanna L. Borders of ponds. Aug. Torrey collection. Shelburne Pond, (Perkins); Perkins' Flora.
- Aster junceus Ait. Swamps and bogs. Aug. Ought to be found here.
- Aster novi-belgii L. Low grounds. Aug., Sept. Torrey collection.
- Bidens vulgata Greene. Has been confused with B. frondosa and should be sought in moist places and by roadsides. Aug., Sept.
- Cirsium discolor (Muhl.) Spreng. Rich soil. Aug., Sept. Ferrisburgh, (Brainerd).
- Hieracium Gronovii L. Sandy soil. Aug., Sept. Burlington, (Torrey); Perkins' Flora.
- **Hieracium marianum** Willd. Open woods and clearings. June-Aug. Torrey collection.
- Lactuca hirsuta Muhl. Dry open ground. July. Charlotte, (Pringle).

- Lactuca integrifolia Bigel. Rich thickets. July, Aug. Reported from Burlington, (Ross).
- Onopordum Acanthium L. Roadsides and waste places. July, Aug. Charlotte, (Pringle). Dry pastures, Williston, (Robbins); Thompson's Vermont.
- Petasites palmatus (Ait.) Gray. Swamps. May, June. Charlotte, (Pringle).
- Prenanthes serpentaria Pursh. Dry open soil. Aug., Sept. Torrey collection.

STATISTICAL SUMMARIES

In the following summaries varieties and hybrids are counted with species. When, however, in Tables II and III the numbers are the same, precedence in the list is given to the family or genus having the fewer varieties and hybrids. The summaries include the species recognized in the list of "Additions and Corrections."

I. NUMBERS AND SYSTEMATIC DISTRIBUTION OF SPECIES

	Native	Foreign	Total
Pteridophytes	67	0	67
Spermatophytes	895	278	1172
Totals	962	278	1240

II. DISTRIBUTION OF SPECIES AND VARIETIES AMONG THE PRINCIPAL FAMILIES

INCLUDING ALL HAVING 15 OR MORE SPECIES

	Genera	Speci	es and varie	ties
		Native	Foreign	Total
Cyperaceae	7	128	0	128
Compositae	37	84	37	121
Gramineae	44	74	37	111
Rosaceae	16	56	9	65
Ranunculaceae	11	31	5	36
Polypodiaceae	15	36	0	36
Orchidaceae	12	34	0	34
Leguminosae	14	13	20	33
Cruciferae	15	15	18	33
Labiatae	18	18	14	32
Ericaceae	16	30	0	30
Polygonaceae	4	16	13	29
Umbelliferae	16	15	6	21
Liliaceae	17	20	4	24
Caryophyllaceae	10	8	16	24
Salicaceae	2	16	6	22
Scrophulariaceae	12	16	5	21
Violaceae	1	18	1	19
Najadaceae	$\frac{2}{7}$	18	0	18
Caprifoliaceae	7	15	2	17
Onagraceae	4	13	$\frac{2}{3}$	16
Urticaceae	10	10	5	15
The other 89 families	168	279	76	355

III. THE PRINCIPAL GENERA

INCLUDING ALL HAVING 10 OR MORE SPECIES

Genera	Species	Genera	Species
Carex	94	Salix	15
Viola	19	Ranunculus	15
Polygonum	18	Scirpus	12
Potamogeton	18	Habenaria	11
Aster	16	Crataegus	11
Panicum	16	Galium	10
Solidago	15	Juneus	10

ADDITIONS AND CORRECTIONS

- Preface. After C. G. Hinsdale, Boston, Mass., insert—A. S. Hitchcock, Washington, D. C.
- Page 14. Insert—Panicum linearifolium Scribn. Woods. June-Aug. Burlington, (Hitchcock).
- Page 16. Third line from bottom, before Hordeum sativum Jessen, insert—Avena sativa L., Oat.
- Page 19. Sixteenth line from bottom, for var. irrigua (Wahlenb.) Fern, read var. pallens Fern.
- Page 20. Twelfth line from top, for "Low open ground; frequent." read "Essex Junction, (Brainerd)."
- Page 35. Tenth line from bottom, before "South Burlington, (N. F. F.)" insert—Burlington, (Torrey).
- Page 45. Omit first, second and third lines. Before DENTARIA insert—
 - CONRINGIA (Heist.) Adans. Hare's-ear Mustard
- C. ORIENTALIS (L.) Dumort. Adventive in railroad yards, Burlington, (N. F. F.). June, July.
- Page 53. Tenth line from bottom, for "(Torrey)" read "(Benedict)."
- Page 65. Seventh line from top, for "Colchester, (Torrey, N. F. F.)," read "Burlington, (Torrey); Colchester, (N. F. F.)."
- Page 65. Twelfth line from bottom, after "Fort Ethan Allen, (Grout)," insert—railroad yard, Burlington, (N. F. F.).
- Page 73. Fourth line from bottom, for "I. COCCINEA L." read "I. COCCINEA L. var. HEDERIFOLIA (L.) Gray."
- Page 80. Ninth line from top, for "Burlington and Essex, (N. F. F.)," read "Burlington, (Torrey, N. F. F.); Essex, (N. F. F.)."
- Page 80. Tenth line from bottom, for "Blue Bottle" read "Bluebottle."
- Page 94. Sixth line from top, after "those" insert—a comma.

Abele 29	Alyssum, Hoary 45	Purple Thorn 78
Abies 6	Sweet 46	Thorn 78
Abutilon 61	Amaranth 37	Aquifoliaceae 58
Acacia, False 55	Family 37	Aquilegia 41
Acalypha 57	Green 37	Arabis 44
Acer 59	Prostrate 37	Araceae 23
Aceraceae 59	Thorny 37	Aralia 66
Achillea 85	Amaranthaceae 37	Araliaceae 66
Acnida 37	Amaranthus 37	Arbor Vitae 7
Acorus 23	Ambrosia 85	Arbutus, Trailing 69
Actaea 40	Amelanchier 49	Arceuthobium 34
Adder's Mouth 29	American Aspen 30	Arctium 86
Green 29	Beech 32	Arctostaphylos 69
Adder's Tongue 4	Bladder Nut 59	Arenaria 38
Family 4	Brooklime 80	Arethusa 27
Yellow 25	Cranberry 71	Arisaema 23
Adiantum 1	Elm 34	Aristida 10
Adlumia 44	Fly Honeysuckle 82	Aristolochiaceae 34
Aesculus 59	Germander 78	Aromatic Winter-
Agastache 75	Hop Hornbeam 32	green 69
Agrimonia 49	Hornbeam 32	Arrhenatherum 10
Agrimony 49	Larch 6	Arrow Grass Family 9
Tall Hairy 49	Mountain Ash 51	Arrow-head 9
Woodland 49	Orpine 48	Arrow-leaved Tear-
Agropyron 9	Pennyroyal 76	thumb 36
Agrostemma 38	White Hellebore 26	Arrow-wood 83
Agrostis 10	Yew 6	Downy 83
Ague-weed 72	Amphicarpa 53	Maple-leaved 83
Aizoaceae 38	Anacardiaceae 58	Toothed 83
Ajuga 75	Anagallis 71	Artemisia 86
Alder 31	Anaphalis 85	Artichoke, Jerusa-
Black 58	Andromeda 69	lem 89
Downy Green 31	Andropogon 10	Arum, Dragon 23
Hoary 31	Anemone 41	Family 23
Smooth 32	Cut-leaved 41	Water 23
Speckled 31	Long-fruited 41	Asarabacca 34
Alder-leaved Buck-	Round-headed 41	Asarum 34
thorn 60	Tall 41	Asclepiadaceae 73
Alfalfa 54	Wood 41	Asclepias 73
Alisma 9	Angelica 66	Ash 72
Alismaceae 9	Antennaria 85	American Moun-
Allium 25	Anthemis 85	tain 51
Allium 25	Anthoxanthum 10	Black 72
Allspice, Wild 43 Alnus 31	Apios 53	European Moun-
Alopecurus 10	Apocynaceae 72	tain 51
Alsike Clover 55	Apocynum 72	Green 72
Alternate-leaved Dog-	Appalachian Cherry 51	Northern Prickly 56
wood 68	Apple 51	Prickly 56
wood oo	of Peru 78	Red 72

White 72	Barren Strawberry 53	
Ash-leaved Maple 59	Bartonia 72	Betula 32
Spiraea 53	Yellow 72	Betulaceae 31
Asparagus 25	Basil 77	Bidens 87
Garden 25	Thyme 77	Bindweed 73
Aspen 29	Basswood 61	Black 35
American 30	Bastard Pennyroyal 78	Field 73
Large-toothed 30	Toad-flax 34	Fringed Black 35
Aspidium 1	Beach Heather 62	Hedge 73
Asplenium 2	Pea 54	Upright 73
Aster 86	Beak Rush 22	Birch 32
Flat-topped White 87		Black 32
Hairy-stemmed 86	Willow 31	Canoe 32
Heart-leaved 86	Bean, Wild 53	Cherry 32
Large-leaved 86	Bearberry 69	Family 32
New England 86	Red 69	Gray 32
Panicled 86	Beard, Goat's 92	Old Field 32
Pringle's 86	Grass 10	Paper 32
Purple-stemmed 87	Bearded Darnel 13	Sweet 32
Small White 87	Beard-tongue 80	White 32
Stiff 86	Bedstraw 81	Yellow 32
Tall White 86	Marsh 82	Bird Cherry 51
Wavy-leaved 87	Northern 82	Birthroot 26
White Wood 86	Rough 81	Birthwort Family 34
Astragalus 53	Small 82	Bishop's Cap 48
Avena 100	Sweet-scented 82	Bitter Cress 45
Avens 50	Beech 32	Dock 36
Purple 50	American 32	Bitternut Hickory 31
Rough 50	Blue 32	Bittersweet 58, 78
Water 50	Family 32	Climbing 58
White 50	Fern 3	Shrubby 58
Yellow 50	Water 32	Bitter-weed 85
Awned Cyperus 21	Beech-drops 81	Black Alder 58
Wheat Grass 9	False 70	Ash 72
Azalea, Pink 70	Beggar's Lice 74	Bindweed 35
Bachelor's Button 87	Beggar's-ticks, Com-	Birch 32
Balm of Gilead 30	mon 87	Chokeberry 51
Fir 6	Swamp 87	Cohosh 41
Balmony 79	Bellflower 84	Dewberry 52
Balsam 60	European 84	Gum 68
Fir 6	Marsh 84	High Blueberry 70
Groundsel 91	Nettle-leaved 84	Huckleberry 69
Poplar 29	Bellwort 26	Larch 6
Balsam-apple, Wild 84	Large-flowered 26	Medick 54
Balsaminaceae 60	Sessile-leaved 26	Mustard 45
Baneberry 40	Benjamin Bush 43	Oak 33
Red 40	Bent Grass 10	Raspberry 52
White 40	Benzoin 43	Snakeroot 41, 67
Barbarea 44	Berberidaceae 43	Spruce 6
Barberry 43	Berberis 43	Sugar Maple 59
Common 43	Bergamot, Wild 76	Willow 30
Family 43	Berry, Partridge 82	Blackberry, High
Barley 13, 16	Pigeon 37	Bush 52
Egyptian 13	Service 49	Recurved 52
Barnyard Grass 12	Berteroa 45	Running Swamp 52

Black-eyed Susan 90	Borage Family 74	Crowfoot 42
Black-fruited Moun-	Boraginaceae 74	Bull Thistle 88
tain Rice 14	Botrychium 4	Bulrush 22
Bladder Campion 39	Bottle, Blue 87	Great 22
Fern 2	Grass 16	River 22
Ketmia 61	Bottle-brush Grass 13	Bunchberry 68
Nut 59	Bouncing Bet 39	Bur Clover 54
Nut Family 59	Box Elder 59	Grass 11
Bladderwort 80	Brachyelytrum 10	Marigold 87
Family 80	Bracken 3	Oak 33
Greater 80	Brake 3	Burdock 86
Blite, Strawberry 37	Cliff 3	Common 86
Bloodroot 43	Purple Cliff 3	Great 86
Blue Beech 32	Rock 2	Burnet 53
Bottle 87	Slender Rock 2	Garden 53
Cohosh 43	Bramble 52	Bur-reed 7
Curls 78	Brasenia 40	Broad-fruited 7
Devil 74	Brassica 45	Family 7
Flag 27	Brier, Cat 26	Burseed 74
Myrtle 73	Green 26	Bush, Benjamin 43
Sailors 87	Bristly Crowfoot 42	Clover 54
Vervain 75	Foxtail Grass 16	Fever 43
Vetch 55	Sarsaparilla 66	Honeysuckle 82
Bluebell 84	Broad Beech Fern 3	Rock Shad 49
Family 84	Broad-fruited Bur-	Shad 49
Bluebells 75	reed 7	Shore Shad 49
Blueberry 70	Broad-leaved Spring	Spice 43
Black High 70	Beauty 40	Steeple 53
Early Sweet 71	Brome Grass 11	Bushy Golden-rod 91
High 70	Bromus 11	Butter-and-Eggs 79
Late Low 71	Brooklime, American 8	0Buttercup 42
Low Sweet 71	Broom Beard Grass 10	Bulbous 42
Sour-top 70	Broom-rape 81	Creeping 42
Swamp 70	Family 81	Swamp 43
77 . 3		
Velvet-leaf 70	Brown Beak Rush 22	Tall 42
Bluets 82		Tall 42 Butternut 31
Bluets 82 Blue-eyed Grass 27	Brown Beak Rush 22	Tall 42 Butternut 31 Butterweed 88
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11	Brown Beak Rush 22 Brownish-fruited	Tall 42 Butternut 31
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden-	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk-	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamus 23 Calamis 77
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamus 23 Calamus 77 California Rose 73
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Cana-	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden-rod 91 Blue-weed 74 Blunt-leaved Milk-weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamus 23 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden-rod 91 Blue-weed 74 Blunt-leaved Milk-weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57 Calopogon 27
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden-rod 91 Blue-weed 74 Blunt-leaved Milk-weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69 Rush 24	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48 Bugbane 41	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaecae 57 Callitriche 57 Calopogon 27 Caltha 41
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69 Rush 24 Spruce 6	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48 Bugbane 41 Bugle Weed 75, 76	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57 Calopogon 27 Caltha 41 Camelina 45
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69 Rush 24 Spruce 6 Willow 30	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48 Bugbane 41 Bugle Weed 75, 76 Bugloss, Viper's 74	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57 Calopogon 27 Caltha 41 Camelina 45 Campanula 84
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden-rod 91 Blue-weed 74 Blunt-leaved Milk-weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69 Rush 24 Spruce 6 Willow 30 Wintergreen 70	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48 Bugbane 41 Bugle Weed 75, 76 Bugloss, Viper's 74 Bulb-bearing Loose-	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57 Calopogon 27 Caltha 41 Camelina 45 Campanula 84 Campanula 84
Bluets 82 Blue-eyed Grass 27 Blue-joint Grass 11 Blue-stemmed Golden- rod 91 Blue-weed 74 Blunt-leaved Milk- weed 73 Sandwort 38 Blunt-lobed Woodsia 3 Boehmeria 33 Bog Club Moss 5 Golden-rod 92 Rosemary 69 Rush 24 Spruce 6 Willow 30	Brown Beak Rush 22 Brownish-fruited Rush 24 Buckbean 72 Buckeye 59 Buckthorn 60 Alder-leaved 60 Common 60 Family 60 Buckwheat 35 Climbing False 36 Family 36 Buffalo-berry, Canadian 64 Buffalo Currant 48 Bugbane 41 Bugle Weed 75, 76 Bugloss, Viper's 74	Tall 42 Butternut 31 Butterweed 88 Buttonbush 81 Buttonwood 49 Cabbage, Skunk 23 Calamus 23 Calamagrostis 11 Calamint 77 California Rose 73 Calla 23 Wild 23 Callitrichaceae 57 Callitriche 57 Calopogon 27 Caltha 41 Camelina 45 Campanula 84

Red 39	Cedar, Red 6	Purple 51
White 38	White 7	Red 51
Camptosorus 2	Celandine 43	Choke Cherry 51
Canada Blue Grass 15	Celastraceae 58	Christmas Fern 3
Golden-rod 91	Celastrus 58	Chryanthemum 87
Hawkweed 89	Celtis 33	Chrysosplenium 48
Plum 51	Cenchrus 11	
		Cicely, Smoother
Thistle 88	Centaurea 87	Sweet 67
Violet 62	Cephalanthus 81	Sweet 67
Canadian Buffalo-	Cerastium 38	Woolly Sweet 67
berry 64	Chain Fern 3	Cicer 53
Canary Grass 15	Chair-maker's Rush 22	Cichorium 87
Cancer-root 81	Chamaedaphne 69	Cicuta 66
One-flowered 81	Chamomile 85	Cimicifuga 41
Canker-root 41	Wild 90	Cinna 11
Cannabis 33	Charlock 45	Cinnamon Fern 3
Canoe Birch 32	Cheat 11	Rose 52
Caper Family 47	Checkerberry 69	Cinquefoil 50
Capillaire 69	Cheeses 61	Marsh 51
Capparidaceae 47	Chelidonium 43	Rough 51
Caprifoliaceae 82	Chelone 79	Shrubby 51
Capsella 45	Chenopodiaceae 36	
		Silvery 50
Caraway 66	Chenopodium 36	Tall 51
Cardamine 45	Cherry 51	Circaea 64
Cardinal Flower 84	Appalachian 51 Birch 32	Cirsium 88
Carduus 87		Cistaceae 62
Carex 17	Bird 51	Clammy Ground Cher-
Carnation 38	Choke 51	ry 78
Carpenter-weed 77	Clammy Ground 78	Hedge Hyssop 79
Carpet Weed 38	Fire 51	Locust 55
Carpinus 32	Ground 78	Clammy-weed 47
Carrion-flower 26	Pin 51	Claytonia 40
Carrot 67	Rum 51	Clearweed 34
Wild 67	Sand 51	Cleavers 81
Carum 66	Wild Black 51	Clematis 41
Carya 31	Wild Red 51	Purple 41
Caryophyllaceae 38	Chess 11	Wild 41
Cashew Family 58	Wild 11	Cliff Brake 3
Cassandra 69	Wood 11	Climbing bitter-sweet 58
Castalia 40	Chestnut 32	False Buckwheat 36
Castanea 32	Oak 33	Fumitory 44
Cat Brier 26	Chick-pea 53	Nightshade 78
Mint 76	Chickweed 39	Clintonia 25
Spruce 6	Common 39	Yellow 25
Catchfly 39	Common Mouse-ear 38	SClinton's Shield Fern 1
Night-flowering 39	Field Mouse-ear 38	Closed Gentian 72
Sleepy 39	Indian 38	Clotbur 92
Sweet William 39	Mouse-ear 38	Clover 55
Catnip 76	Nodding 38	Alsike 55
Cat-tail, Common 7	Wintergreen 72	Bur 54
Family 7	Chicory 87	Bush 54
Flag 7	Common 87	Нор 55
Narrow-leaved 7	Chimaphila 69	Low Hop 55
Caulophyllum 43	Chiogenes 69	Rabbit-foot 55
Ceanothus 60	Chokeberry, Black 51	Red 55
	The state of the s	

O4 FF	C1 C0	III. 1 10
Stone 55	Cornus 68	Water 46
Sweet 54	Corpse Plant 70	Winter 44
White 55	Corydalis 44	Yellow 46
White Sweet 54	Golden 44	Crested Shield Fern 1
Yellow 55	Pale 44	Crowfoot 42
Yellow Sweet 54	Pink 44	Bristly 42
Zigzag 55	Corylus 32	Bulbous 42
Club Moss 5	Cotton Grass 22	Common White
Moss Family 5	Cotton-wood 30	Water 42
Rush 22	Couch Grass 9	Cursed 42
Clustered Snakeroot 6		Family 40
Coast Jointweed 35		
	Lily 40	Hooked 42
Cockle, Corn 38	Parsnip 67	Small-flowered 42
Cocklebur 92	Wheat 79	Stiff Water 42
Coffee, Wild 83	Cowbane, Spotted 66	Tall 42
Cohosh 40	Cow-herb 39	Yellow Water 42
Black 41	Cowslip, English 71	Cruciferae 44
Blue 43	Virginian 77	
	Virginian 75	Cryptogramma 2
Coltsfoot 92	Cowslips 41	Cryptotaenia 66
Columbine 41	Crack Willow 30	Cucumber, Wild 84
Garden 41	Cranberry 70	One-seeded Bur 84
Wild 41	American 71	Star 84
Comandra 34	High-bush 83	Cucumber-root, In-
Comfrey 75		diam 05
Common 75	Large 71	dian 25
	Small 71	Cucurbitaceae 84
Wild 74	Tree 83	Cudweed 89
Compositae 85	Cranesbill 56	Low 89
Composite Family 85	Wild 56	Cup Plant 91
Cone-flower 90	Crassulaceae 47	Huntman's 47
Tall 90	Crataegus 49	Currant 48
Conium 66		
	Cream-colored Vetch-	Buffalo 48
Convallaria 25	ling 54	Garden Red 48
Convolvulaceae 73	Creeper, Disk 60	Missouri 48
Convolvulvus 73	Tendril 60	Swamp Red 48
Family 73	Virginia 60	Wild Black 48
Coptis 41	Creeping Buttercup 42	Cursed Crowfoot 42
Coral Root 27	Eragrostis 12	
Early 27		Cuscuta 73
	Snowberry 69	Cut-grass 13
Large 27	Spearwort 42	Rice 13
Corallorrhiza 27	Cress, Bitter 45	Cut-leaved Anemone 41
Cord Grass 16	Common Winter 45	Bugle Weed 76
Cork Elm 34	Cow 46	Golden-rod 91
Corn Cockle 38	Erect-fruited Win-	Pepper-root 45
Flower 87	ter 44	
Indian 16		Toothwort 45
	Field 46	Cynoglossum 74
Poppy 43	Field Penny 47	Cyperaceae 17
Speedwell 80	Hairy Marsh 46	Cyperus 21
Spurrey 39	Hairy Rock 44	Awned 21
Squirrel 44	Lake 46	Shining 21
Wild Broom 15	Marsh 46	
Cornaceae 68	Penny 47	Straw-colored 21
Cornel 68		Cypress Spurge 57
Dwarf 68	Purple Rock 44	Cypripedium 27
	Rock 44	Cystopteris 2
Round-leaved 68	Small Bitter 45	Dactylis 11
Silky 68	Smooth Rock 44	Daisy Fleabane 88

109

Michaelmas 87	Red-osier 68	Eleocharis 21
Ox-eye 87	Round-leaved 68	Elm 34
White 87	Silky 68	American 34
Yellow 90	Dog's-tooth Violet 25	Cork 34
Dalibarda 50	Doll's Eyes 40	English 34
Dandelion 92	Doorweed 35	Red 34
Common 92	Downy Arrow-wood 83	Rock 34
Red-seeded 92	Golden-rod 92	Slippery 34
Danthonia 11	Green Alder 31	White 34
Daphne 64	Rattlesnake Plan-	Elodea 9
Darnel 13	tain 28	Elymus 12
Bearded 13	Yellow Violet 63	Enchanter's Night-
Common 13	Draba 46	shade 64
Datura 78	Dragon Arum 23	English Cowslip 71
Daucus 67	Drop-seed 14, 16	Elm 34
Day Lily 25	Dropwort 50	Hawthorn 49
Dead Nettle 76	Drosera 47	Plantain 81
Decodon 64	Droseraceae 47	Epifagus 81
Dentaria 45	Duck's-meat .23	Epigaea 69
Deschampsia 11	Duckweed 23	Epilobium 64
Desmodium 53	Family 23	Epipactis 28
Devil's Paint-brush 89	Greater 23	Equisetaceae 4 Equisetum 4
Dewberry, Black 52	Ivy-leaved 23	Equisetum 4 Eragrostis 12
Red 52	Lesser 23	
Swamp 52	Dulichium 21	Creeping 12 4 Strong-scented 12
Dianthus 38	Dutchman's Breeches 4 Dwarf Cornel 68	Erechtites 88
Dicentra 44		Erect-fruited Winter
Dicksonia 2 Diervilla 82	Ginseng 66 Mistletoe 34	Cress 44
Digitaria 12	Raspberry 52	Erect Knotweed 35
Direa 64	Sumach 58	Ericaceae 69
Disk Creeper 60	Dyssodia 88	Erigeron 88
Ditch Stonecrop 47	Early Coral Root 27	Eriocaulaceae 23
Dock 36	Golden-rod 91	Eriocaulon 23
Bitter 36	Meadow Parsnip 68	Eriophorum 22
Great Water 36	Meadow Rue 43	Erodium 56
Patience 36	Saxifrage 48	Eryngium 67
Spatter 40	Sweet Blueberry 71	Eryngo 67
Swamp 36	Yellow Violet 63	Erysimum 46
Yellow 36	Ebony Spleenwort 2	Erythronium 25
Dock-leaved Persi-	Echinochloa 12	Eupatorium 88
caria 35	Echinocystis 84	Euphorbia 57
Dockmackie 83	Echium 74	Euphorbiaceae 57
Dodder 73	Eel Grass 9	European Bellflower 84
Dog Fennel 85	Eglantine 52	Mallow 60
Violet 62	Egyptian Barley 13	Millet 14
Dogbane 72	Elaeagnaceae 64	Mountain Ash 51
Family 72	Elder 83	Stickseed 74
Spreading 72	Box 59	Evening Primrose 65
Dogberry 48	Common 83	Primrose Family 64
Dogwood 68	Poison 58	Evergreen Wood Fern 1
Alternate-leaved 68	Red-berried 83	Everlasting 85
Family 68	Wild 66	Common 89
Panicled 68	Elderberry 83	Pea 54
Poison 58	Elecampane 90	Pearly 85

	Tri state Obain 9	0:3 331- 47
Plantain-leaved 85	Virginia Chain 3	Side-saddle 47
Fagaceae 32	Wood 1	Star 72
Fagopyrum 35	Fescue Grass 12	Wind 41
Fagus 32	Meadow 12	Flowering Fern 3
False Acacia 55	Nodding 12	Fern Family 3
Beech Drops 70	Sheep's 12	Sage 76
Dragon Head 77	Red 12	Spurge 57
Flax 45	Taller 12	Wintergreen 57 Flower-of-an-hour 61
Hellebore 26	Festuca 12	Foam-flower 48
Loosestrife 65	Fetid Marigold 88	Forget-me-not 75
Mermaid 58	Fever Bush 43	True 75
Mermaid Family 58	Feverwort 83 Field Bindweed 73	Forked Beard Grass 10
Miterwort 48	Cress 46	Four-leaved Milkweed 73
Nettle 33	Golden-rod 91	Fowl Meadow Grass 15
Pimpernel 79		Foxtail Grass 10
Solomon's Seal 26	Milkwort 57	Green 16
Spikenard 26	Mouse-ear Chick-	Yellow 16
Feather Geranium 36	weed 38	
Prince's 35	Penny Cress 47	Fragaria 50 Fragile Bladder Fern 2
Fennel, Dog 85	Sorrel 36	Fraxinus 72
Fen Orchis 29	Sow Thistle 92	Fringed Black Bind-
Fern, Beech 3	Spurrey 39	weed 35
Bladder 2	Figwort 80	Loosestrife 71
Boott's Shield 1	Family 79	Orchis 28
Broad Beech 3	Filbert 32	
Chain 3	Filipendula 50	Polygala 57
Christmas 3	Finger Grass 12	Frog's Bit Family 9 Frost Grape 60
Cinnamon 3	Fiorin 10 Fir 6	Frostweed 62
Clinton's Shield 1	Balm-of-Gilead 6	Fumaria 44
Common Bladder 2	Balsam 6	Fumariaceae 44
Crested Shield 1	Fire Cherry 51	Fumitory 44
Evergreen Wood 1	Fireweed 65, 88	Climbing 44
Family 1	Five-finger 50, 51	Common 44
Flowering 3	Flag, Blue 27	Family 44
Fragile Bladder 2	Cat-tail 7	Galeopsis 75
Goldie's 1	Sweet 23	Galingale 21
Grape 4	Flat-topped White As-	Galinsoga 88
Hay-scented 2	ter 87	Galium 81
Interrupted 4	Flax 55	Gall-of-the-earth 90
Lady 2		Garden Asparagus 25
Lance-leaved Grape 4	False 45	Burnet 53
Long Beech 3	Family 55	Columbine 41
Marginal Shield 1	Fleabane 88	Heliotrope 84
Marsh 1	Daisy 88	Orpine 47
New York 1	Pink 88	Phlox 74
Oak 3	Fleur-de-lis 27	Red Currant 48
Ostrich 2	Floating Foxtail	Snowberry 83
Rattlesnake 4	Grass 10	Sorrel 36
Royal 4	Manna Grass 13	Garget 37
Sensitive 2	Floerkea 58	Garlie 25
Shield 1	Flower, Cardinal 84	Gaultheria 69
Spinulose Shield 1	Corn 87	Gaylussacia 69
Sweet 31	Moccasin 27	Gentian 72
Ternate Grape 4	Monkey 79	Closed 72

	INDEX	111
Family 72	Gooseberry 48	Lyme 12
Horse 83	Prickly 48	Manna 13
Stiff 72	Goosefoot 36	Marsh 16
Gentiana 72	Family 36	Meadow 15
Gentianaceae 72	Maple-leaved 37	Meadow Foxtail 10
Geraniaceae 56	Oak-leaved 37	Melic 13
Geranium 56	Upright 37	Millet 16
Family 56	Gourd Family 84	Oat 10
Feather 36	Gramineae 9	of Parnassus 48
Gerardia 79	Grape 60	Old-witch 14
Slender 79	Fern 4	Orchard 11
Small-flowered 79	Frost 60	Pale Manna 13
Germander 78	River-bank 60	Panic 14
American 78	Summer 60	Perennial Ray 13
German Millet 16	Grass	Pigeon 16
Geum 50	Awned Wheat 9	Pink 27
Giant Hyssop 75	Barnyard 12	Poverty 10
Gilia 74	Beard 10	Quick 9
Gill-over-the-ground 77	Bent 10	Quitch 9
Ginger, Wild 34	Blue-eyed 27	Rattlesnake 13
Ginseng 66	Blue-joint 11	Reed Bent 11
Dwarf 66	Bottle 16	Reed Canary 15
Family 66	Bottle-brush 13	Reed Meadow 13
Glaucous Honey-	Bristly Foxtail 16	Rib 81
suckle 83	Brome 11	Ribbon 15
Willow 30	Broom Beard 10	Ripple 81
Glyceria 13	Bur 11	Rough Foxtail 16
Gnaphalium 89	Canada Blue 15	Rush 16
Goat's Beard 92	Canary 15	Rye 13
Golden Alexanders 68	Common Hair 11	Seneca 13
Corydalis 44	Common Wild Oat 11	Slender Wheat 9
Hedge Hyssop 79	Cord 16	Slender Wood Reed 11
Meadow Parsnip 68	Cotton 22	Slough 16
Ragwort 91	Couch 9	Small Crab 12
Saxifrage 48	Eel 9	Spear 15
Seal 42	Family 9	Squirrel-tail 13
Golden-rod 91	Fescue 12	Sweet Vernal 10
Blue-stemmed 91	Finger 12	Tall Oat 10
Bog 92	Floating Foxtail 10	Tape 9
Bushy 91	Floating Manna 13	Thin 10
Canada 91	Forked Beard 10	Triple-awned 10
Cut-leaved 91	Fowl Meadow 15	Vanilla 13
Downy 92	Foxtail 10	Water Star 24
Early 91	Goose 81	Weak Spear 15
Field 91	Grove Spear 15	White 13
Gray 91	Hair 10 Herd's 15	White Bent 10 Wild Oat 11
Hairy 91 Late 92	Holy 13	Wire 15
Stout Ragged 92	Hungarian 16	Wood 16
Tall Hairy 92	Hungarian Brome 11	Wood Reed 11
White 91	Indian 16	Wood Reed II
Zigzag 91	June 15	Yellow Nut 21
Goldie's Fern 1		Gratiola 79
Goldthread 41		Graveyard Moss 57
Goose Grass 81		Gray Birch 32
		•

Golden-rod 91	Panicled 89	Alyssum 45
Oak 33	Rough 90	Hobble-bush 83
Great Bulrush 22	Hawthorn 49	Hog Peanut 53
Burdock 86	English 49	Hog-weed 85
Ragweed 85	Hay-scented Fern 2	Holly 58
St. John's-wort 61	Hazelnut 32	Family 58
	Beaked 32	Mountain 58
Water Dock 36		Holy Grass 13
Willow-herb 65	Heal-all 77	Honewort 66
Greater Bladderwort 80		
Duckweed 23	Twayblade 29	Honeysuckle 82
Green Adder's Mouth 29		American Fly 82
Amaranth 37	Heart's-ease 62, 63	Bush 82
Ash 72	Heath Family 69	Family 82
Brier 26	Heather, Beach 62	Glaucous 83
Foxtail 16	Hedeoma 76	Tartarian 83
Greenish-flowered Win-		Hooked Crowfoot 42
tergreen 70	Hyssop 79	Hop 33
Grim the Collier 89	Mustard 46	Clover 55
Gromwell 74	Nettle 77	Common 33
Common 75	Helianthemum 62	Hornbeam 32
Ground Cherry 78	Helianthus 89	Japanese 33
Hemlock 6	Heliopsis 89	Hordeum 13, 16
Ivy 77	Heliotrope, Garden 84	Horehound, Water 76
Laurel 69	Hellebore, American	Hornbeam 32
Pine 5	White 26	American 32
Pink 73	False 26	American Hop 32
Ground-nut 53	Hemerocallis 25	Hop 32
Groundsel 91	Hemlock 7	Horse Gentian 83
Balsam 91	Ground 6	Mint 76
Common 91	Poison 66	Horse-chestnut 59
Grove Spear Grass 15	Water 66	Common 59
Gum, Black 68	Hemp 33	Horseradish 46
Sour 68	Common 33	Horsetail 4
Gypsophila 38	Indian 72	Common 4
Low 38	Nettle 75	Family 4
Habenaria 28	Water 37	Marsh 5
Hackberry 33	Henbit 76	Shore 5
Hackmatack 6	Hepatica 41	Wood 5
Hair Grass 10	Heracleum 67	Horseweed 88, 90
Hairy Golden-rod 91	Herb Robert 56	Hound's Tongue 74
Marsh Cress 46	Herd's Grass 15	Common 74
Rock Cress 44	Heteranthera 24	Houstonia 82
Wood Rush 24	Hibiscus 61	Huckleberry 69
Hairy-stemmed Aster 8	6Hickory 31	Black 69
Halberd-leaved Tear-	Bitternut 31	Hudsonia 62
thumb 35	Shagbark 31	Humulus 33
Haloragidaceae 65	Shellbark 31	Hungarian Brome
Hamamelidaceae 49	Swamp 31	Grass 11
Hamamelis 49	White-heart 31	Grass 16
Hardhack 53	Hieracium 89	Huntsman's Cup 47
Harebell 84	Hierochloe 13	Husk Tomato 78
Hare's Tail 22	High Blueberry 70	Hydrastis 42
Hawkweed 89	High-bush Blackberry 5	
Canada 89	Cranberry 83	Hydrocotyle 67
Orange 89	Hoary Alder 31	Hydrophyllaceae 74
	ary much of	11, drophy naceae 14

TI-duanhadhum 74	Cyang 15	Yellow Pond Lily 40
Hydrophyllum 74.	Grass 15	Large-bracted Plan-
Hypericaceae 61.	Pink 70 Juniper 6	tain 81
Hypericum 61	Common 6	Large-flowered Bell-
Hyssop, Clammy		wort 26
Hedge 79	Juniperus 6	Large-fruited Snake-
Giant 75	Kalmia 69	
Golden Hedge 79	Kentucky Blue Grass 1 King Devil 89	Large-leaved Aster 86
Hedge 79	Kinnikinnik 68	Orchis 28
Hystrix 13	Knotgrass 35	Pondweed 8
Ilex 58	Knotted Rush 24	Large-toothed Aspen 30
Ilysanthes 79	Knotted Rush 24 Knotweed 35	Larix 6
Impatiens 60 Indian Chickweed 38	Erect 35	Late Golden-rod 92
Corn 16	Japanese 35	Low Blueberry 71
Cucumber-root 25	Labiatae 75	Lathyrus 54
Grass 16	Labrador Tea 69	Lauraceae 43
	Lactuca 90	Laurel 69
Hemp 72	Ladies' Tobacco 85	Family 43
Mallow 61 Mustard 45	Ladies' Tresses 29	Ground 69
	Nodding 29	Pale 69
Pipe 70 Poke 26	Slender 29	Sheep 69
Rice 16	Lady Fern 2	Laurestinus 83
Tobacco 84	Lady's Slipper 27	Leaf, Leather 69
Turnip 23	Large Yellow 27	Velvet 61
Innocence 82	Pink 27	Leather Leaf 69
Interrupted Fern 4	Ram's Head 27	Leatherwood 64
Inula 90	Showy 27	Lechea 62
Ipomoea 73	Smaller Yellow 27	Ledum 69
Iridaceae 27	Stemless 27	Leek, Wild 25
Iris 27	Lady's Sorrel 56	Leersia 13
Family 27	Thumb 36	Leguminosae 53
Wild 27	Lake Cress 46	Lemna 23
Yellow 27	Lambkill 69	Lemnaceae 23
Ironwood 32	Lamb's Quarters 36	Lentibulariaceae 80
Ivy, Ground 77	Lamium 76	Leonurus 76
Poison 57	Lance-leaved Grape	Lepidium 46
Ivy-leaved Duckweed		Lespedeza 54
Jack-in-the-Pulpit 23	Loosestrife 71	Lesser Duckweed 23
Jamestown Weed 78	Violet 63	Rattlesnake Plan-
Japanese Hop 33	Laportea 33	tain 28
Knotweed 35	Lappula 74	Stitchwort 39
Jerusalem Artichoke	89Larch 6	Lettuce 90
Oak 36	American 6	Prickly 90
Jewelweed 60	Black 6	Tall Blue 90
Jimson Weed 78	Large Coral Root 27	Tall White 90
Joe-Pye Weed 88	Crab Grass 12	White 90
Jointed Rush 24	Cranberry 71	Wild 90
Jointweed, Coast 35	Pepper-root 45	Leverwood 32
Juglandaceae 31	Purple-fringed Or-	Levisticum 67
Juglans 31	chis 28	Lilac 72
Juncaceae 24	Toothwort 45	Common 72
Juncaginaceae 9	Twayblade 28	Liliaceae 25
Juneus 24	Yellow Lady's Slip-	Lilium 25
June Berry 49	per 27	Lily 25
-		

Common Day 25	Loose-spiked Milk-	Malvaceae 61
Cow 40	wort 57	Manna Grass 13
Day 25	Loosestrife 64, 71	Maple 59
Family 25	Bulb-bearing 71	Ash-leaved 59
Large Yellow Pond 4		Black Sugar 59
Meadow 25	Family 64	Family 59
of the Valley 25	Fringed 71	Mountain 59
Small Yellow Pond 4		Red 59
Sweet-scented White	e Swamp 64	Rock 59
	Tufted 71	Silver 59
Water 40		
Tiger 25	Whorled 71	Striped 59
Tuberous White	Lopseed 81	Sugar 59
Water 40	Family 81	Swamp 59
Water 40	Loranthaceae 34	White 59
Wild Orange-red 25	Lousewort 79	Maple-leaved Arrow-
Wild Yellow 25	Common 79	wood 83
Wood 25	Lovage 67	Goosefoot 37
Yellow Pond 40	Love Vine 73	Marginal Shield Fern 1
Limnanthaceae 57	Low Cudweed 89	Marguerite 87
Linaceae 55	Gypsophila 38	Marigold, Bur 87
Linaria 79	Hop Clover 55	Fetid 88
Linden 61	Spear Grass 15	Marsh 41
Family 61	Sweet Blueberry 71	Water 87
Linnaea 82	Lucerne 54	Marsh Bedstraw 82
Linum 55	Ludvigia 65	Bellflower 84
Lion's Heart 77	Lungwort 75	Blue Violet 62
Liparis 28	Lupine 54	Cinquefoil 51
Liquorice, Wild 82	Wild 54	Cress 46
Listera 29	Lupinus 54	Fern 1
Lithospermum 74	Luzula 24	Grass 16
Little Grape Fern 4	Lychnis 38	Horsetail 5
Live-for-ever 47	Lycopodiaceae 5	Marigold 41
Liverleaf 41	Lycopodium 5	Muhlenbergia 14
Round-lobed 42	Lycopus 76	Pea 54
Sharp-lobed 41	Lycium 78	St. John's-wort 62
Lobelia 84	Lyme Grass 12	Skullcap 77
Family 84	Lysimachia 71	Speedwell 80
Lobeliaceae 84	Lythraceae 64	Masterwort 66, 67
Lobularia 46	Lythrum 64	Matricaria 90
Locust 55	Madder Family 81	Matrimony Vine 78
Clammy 55	Wild 82	Mayflower 69
Common 55	Mad-dog Skullcap 77	Mayweed 85
Lolium 13	Maianthemum 25	Meadow Fescue 12
Lombardy Poplar 30	Maidenhair 1	Foxtail Grass 10
Long Beech Fern 3	Spleenwort 2	Grass 15
Long-bracted Orchis 28		Lily 25
Long-fruited Anem-	Common 61	Muhlenbergia 14
one 41	European 61	Queen of the 53
Long-leaved Stitch-	Family 61	Rue 43
wort 39	Indian 61	Meadow-sweet 53
Sundew 47	Musk 61	Medeola 25
Long-spurred Violet 6		Medicago 54
Lonicera 82	Whorled 61	Medick 54
	Malva 61	Black 54
	1.141.44 01	DIACK 94

	1110211	110
Spotted 54	False 48	Mustard 45
Melampyrum 79	Naked 48	Black 45
Melic Grass 13	Two-leaved 48	Common Hedge 47
Melica 13	Mithridate Mustard 47	Family 44
Melilot 54	Moccasin Flower 27	Hedge 46
		Indian 45
White 54	Mock Pennyroyal 76	Mithridate 47
Yellow 54	Mocker Nut 31	
Melilotus 54	Mollugo 38	Tower 44 Treacle 46
Mentha 76	Monarda 76	Tumble 46
Menyanthes 72	Moneses 70	Worm-seed 46
Mercury, Three-	Moneywort 71	
seeded 57	Monkey Flower 79	Myosotis 75
Mertensia 75	Monotropa 70	Myrica 31
Mexican Tea 36	Moonwort 4	Myricaceae 31
Mezereum 64	Moosewood 59, 64, 83	Myriophyllum 65
Family 64	Morning Glory 73	Myrtle, Blue 73
Michaelmas Daisy 87	Common 73	Yellow 71
Microstylis 29	Small Red 73	Naiad 8
Milfoil 85	Wild 73	Najadaceae 8
Water 65	Morus 33	Najas 8
Milk Purslane 57	Moss, Bog Club 5	Naked Miterwort 48
Vetch 53	Club 5	Nannyberry 83
Milkweed 73	Common Club 5	Narrow False Oat 16
Blunt-leaved 73	Common Tree Club	
Common 73	Graveyard 57	tail 7
Family 73	Pink 74	Mountain Mint 77
Four-leaved 73	Shining Club 5	Spleenwort 2
Poke 73	Stiff Club 5	Necklace Poplar 30
Swamp 73	Tree Club 5	Neckweed 80
Milkwort 57	Mossy Stonecrop 47	Nemopanthus 58
Family 57	Mossy-cup Oak 33	Nepeta 76
Field 57	Moth Mullein 80	Nettle 34
Loose-spiked 57	Motherwort 76	Common Hemp 75
Purple 57	Common 76	Dead 76
Racemed 57	Mountain Fringe 44	False 33
Millet 16	Holly 58	Family 33
European 14	Maple 59	Hedge 77
German 16	Mint 77	Hemp 75
Mimulus 79	Rice 14	Rough Hedge 77
Mint 76	Mouse-ear Chickweed 3	
Cat 76	Moxie Plum 69	Spotted Dead 76
Family 75	Mud Plantain 24	Tall Wild 34
Horse 76	Mugwort, Common 86	Tree 33
Mountain 77	Muhlenbergia 14	Wood 33
Narrow-leaved Mount		Nettle-leaved Bell-
tain 77	Meadow 14	flower 84
Spotted 76	Slender 14	New England Aster 86
Whorled 76	Mulberry 33	New Jersey Tea 60
Wild 76	White 33	New York Fern 1
Missouri Currant 48	Mullein 80	Nicandra 78
Mistletoe, Dwarf 34	Common 80	Nigger-head 90
Family 34	Moth 80	Night-flowering Catch-
Mitchella 82	Musk Mallow 61	fly 39
Mitella 48	Storksbill 56	Nightshade 78
Miterwort 48	Musquash Root 66	Climbing 78

Common 79	Pyrola 70	St. John's-wort 61
Enchanter's 64	One-seeded Bur Cucum	
Family 78	ber 84	Pale-leaved Wood Sun-
Smaller Enchant-	One-sided Winter-	flower 89
er's 64	green 70	Panax 66
Nimble Will 14	Onion 25	Panic Grass 14
Nodding Chickweed 38		Panicled Aster 86
Fescue 12	Onoclea 2	Dogwood 68
Ladies' Tresses 29	Ophioglossaceae 4	Hawkweed 89
Trillium 26	Ophioglossum 4	Panicum 14
Wild Rye 12	Orange Hawkweed 89	Pansy 63
Nonesuch 54	Orange-root 42	Papaver 43
Northern Bedstraw 82	Orchard Grass 11	Papaveraceae 43
Prickly Ash 56	Orchidaceae 27	Paper Birch 32
Stitchwort 39	Orchis 29	Pappoose Root 43
Willow-herb 64	Family 27	Parietaria 33
Norway Pine 7	Fen 29	Parnassia 48
Nut, American Blad-	Fringed 28	Parsley Family 66
der 59	Large-leaved 28	Parsnip 67
Bladder 59	Large Purple Fringe	
Mocker 31	28	Early Meadow 68
Nymphaea 40	Long-bracted 28	Golden Meadow 68
Nymphaeaceae 40	Ragged Fringed 28	Poison 67
Nyssa 68	Rein 28	Water 68
Oak 32	Round-leaved 28	Wild 67
Black 33	Showy 29	Partridge Berry 82
Bur 33	Smaller Purple	Pastinaca 67
Chestnut 33	Fringed 28	Pasture Rose 52
Fern 3	Small Wood 28	Patience Dock 36
Gray 33	Tall Green 28	Pea, Beach 54
Jerusalem 36	White Bog 28	Everlasting 54
Mossy-cup 33	White Fringed 28	Marsh 54
Over-cup 33	Orobanchaceae 81	Peanut, Hog 53
Poison 58	Orobanche 81	Pearly Everlasting 85
Red 33	Orpine 47	Pedicularis 79
Swamp White 32	American 48	Pellaea 3
White 32	Family 47	Pellitory 33
Yellow 33	Garden 47	Penny Cress 47
Yellow-barked 33	Oryzopsis 14	Pennyroyal 76
Oakesia 26	Osier 30	American 76
Oak-leaved Goosefoot 37	Osmorhiza 67	Bastard 78
Oat 100	Osmunda 3	Mock 76
Grass 10	Osmundaceae 3	Pennywort, Water 67
Narrow False 16	Ostrich Fern 2	Penthorum 47
Purple 13	Ostrya 32	Pentstemon 80
Oats, Water 16	Over-cup Oak 33	Pepper, Water 35
Oenothera 65	Oxalidaceae 56	Peppergrass 46
Old Field Birch 32	Oxalis 56	Wild 46
Old-witch Grass 14	Ox-eye 89	Pepperidge 68
Oleaceae 72	Daisy 87	Peppermint 76
Oleaster Family 64	Painted Trillium 26	Pepper-root 45
Olive Family 72	Pale Corydalis 44	Cut-leaved 45
Onagraceae 64	Laurel 69	Large 45
One-flowered Cancer-	Manna Grass 13	Two-leaved 45
root 81	Persicaria 35	Pepperwort 46

117

Perennial Ray Grass 1	3 Moss 73	Pokeweed 37
Periwinkle 73	Persicaria 36	Family 37
Common 73	Swamp 70	Polanisia 47
Persicaria, Common 36	Pinus 7	Polemoniaceae 74
Dock-leaved 35	Pinweed 62	Polemonium Family 74
Pale 35	Pipes 4	Polygala 57
Pink 36	Pipewort 23	Fringed 57
Swamp 35	Family 23	Polygalaceae 57
Phalaris 15	Pipsissewa 69	Polygonaceae 35
Phegopteris 3	Pitcher-plant 47	Polygonatum 26
Phleum 15	Family 47	Polygonella 35
Phlox 74	Pitch Pine 7	Polygonum 35
Garden 74	Plane Tree 49	Polypodiaceae 1
Phragmites 15	Family 49	Polypodium 3
Phryma 81	Plant, Corpse 70	Polypody 3
Phrymaceae 81	Cup 91	Polystichum 3
Physalis 78	Plantaginaceae 81	Pondweed 8
Physostegia 76	Plantago 81	Common 8
Phytolacca 37	Plantain 81	Family 8
Phytolaccaceae 37	Common 81	Large-leaved 8
Picea 6	Downy Rattlesnake 2	28Pontederia 24
Pickerel-weed 24	English 81	Pontederiaceae 24
Family 24	Family 81	Poor Man's Weather-
Pigeon Berry 37	Large-bracted 81	glass 71 ·
Grass 16	Lesser Rattlesnake	28 Robin's Plantain 90
Pigweed 36, 37	Mud 24	Poplar 29
Pilea 34	Poor Robin's 90	Balsam 29
Pimbina 83	Rattlesnake 28	Lombardy 30
Pimpernel 71 ·	Robin's 88	Necklace 30
Common 71	Water 9	Silver-leaved 29
False 79	Plantain-leaved Ever-	Trembling 30
Short-stalked False 7	79 lasting 85	White 29
Yellow 68	Platanaceae 49	Poppy 43
Pin Cherry 51	Platanus 49	Corn 43
Pinaceae 6	Plum 51	Family 43
Pine 7	Canada 51	Populus 29
Drops 70	Moxie 69	Portulaca 40
Family 6	Wild 51	Portulacaceae 40
Ground 5	Plumed Thistle 88	Potamogeton 8
Norway 7	Plumeless Thistle 87	Potentilla 50
Pitch 7	Poa 15	Poverty Grass 10
Prince's 69	Pogonia 29	Prairie Willow 30
Red 7	Rose 29	Prenanthes 90
White 7	Small Whorled 29	Prickly Ash 56
Pineapple-weed 90	Whorled 29	Gooseberry 48
Pinesap 70	Poison Dogwood 58	Lettuce 90
Pink 38	Elder 58	Rose 52
Azalea 70	Hemlock 66	Primrose 71
Corydalis 44	Ivy 58	Common Evening 65
Family 38	Oak 58	Evening 65
Fleabane 88	Parsnip 67	Family 71
Grass 27	Sumach 58	Sinuate-leaved Even-
Ground 73	Poke, Common 37	ing 65
June 70	Indian 26	Small-flowered Even-
Lady's Slipper 27	Milkweed 73	ing 65

Primula 71	Ragweed 85	Ribes 48
Primulaceae 71	Great 85	Ribwort 81
Prince's Feather 35	Ragwort 91	Rice Cut Grass 13
Pine 69	Golden 91	Black-fruited Moun-
Pringle's Aster 86	Raisin, Wild 83	tain 14
Prostrate Amaranth 37	Ram's Head Lady's	Indian 16
Prunella 77	Slipper 27	Mountain 14
Prunus 51	Ramsted 79	Slender Mountain 14
Psedera 60	Ranunculaceae 40	Water 16
Pteris 3	Ranunculus 42	White-grained Moun-
Pterospora 70		tain 14
Puccoon 74	Raspberry, Black 52 Dwarf 52	Wild 16
Yellow 42		Richweed 34
	Purple 52	
Pulse Family 53	Purple Flowering 52	
Purple Avens 50	Running 52	River-bank Grape 60
Chokeberry 51	Wild Red 52	River Bulrush 22
Clematis 41	Rattlesnake Fern 4	Robbin's Squaw-weed 91
Cliff Brake 3	Grass 13	Robinia 55
Flowering Rasp-	Plantain 28	Robin's Plantain 88
berry 52	Rattlesnake-root 90	Rock Brake 2
Milkwort 57	Rattlesnake-weed 90	Cress 44
Oat 13	Recurved Blackberry 5	
Raspberry 52	Red Ash 72	Maple 59
Rock Cress 44	Baneberry 40	Sandwort 38
Thorn Apple 78	Bearberry 69	Selaginella 6
Trillium 26	Campion 39	Shadbush 49
Willow 30	Cedar 6	Rocket, Yellow 45
Purple-leaved Willow- herb 65	Chokeberry 51	Rockrose 62
Purple-stemmed As-	Clover 55	Family 62
ter 87	Dewberry 52	Roman Wormwood 85
Purslane 40	Elm 34	Rosa 52
Common 40	Fescue 12	Rosaceae 49
Family 40	Maple 59	Rose 52
Milk 57	Oak 33 Pine 7	California 73
Speedwell 80		Cinnamon 52
Water 65	Spruce 7	Family 49
Pusley 40	Top 10	Mallow 61
Pussy's Toes 85	Red-berried Elder 83	Pasture 52
Pussy Willow 30	Red-osier Dogwood 68 Red-root 60	Pogonia 29
Pycnanthemum 77		Prickly 52
Pyrola 70	Smaller 60	Smooth 52
One-flowered 70	Red-seeded Dande- lion 92	Swamp 52
Pyrus 51	Reed 15	Rosemary, Bog 68
Queen Anne's Lace 67		Rosin-weed 91
Queen of the Meadow 5	Bent Grass 11	Rough Avens 50
Quercitron 33	-	Bedstraw 81
Quercus 32	Meadow Grass 13	Cinquefoil 51
Quick Grass 9	Rein Orchis 28 Rhamnaceae 60	Foxtail Grass 16
Quitch Grass 9	Rhamnus 60	Hawkweed 90
Rabbit-foot Clover 55	Rhododendron 70	Hedge Nettle 77
Racemed Milkwort 57	Rhodora 70	Sunflower 89
Radicula 46	Rhus 58	Round-headed Anem- one 41
Ragged Fringed Or-	Rib Grass 81	Round-leaved Cornel 68
chis 28	Ribbon Grass 15	Dogwood 68
01110 20	remondi Glass 19	Dogwood 08

Orchis 28	Sainfoin 54	Secale 16
Squaw-weed 91		
Sundew 47	St. John's-wort 61	Sedge 17
	Common 62	Family 17
Violet 63	Family 61	Sedum 47
Wintergreen 70	Great 61	Selaginella 6
Round-lobed Liver-	Marsh 62	Rock 6
° leaf 42	Pale 61	Selaginellaceae 6
Rowan Tree 51	Small-flowered 62	Self-heal 77
Royal Fern 4	Spotted 62	Seneca Grass 13
Rubiaceae 81	Salicaceae 29	Snakeroot 57
Rubus 52	Salix 30	Senecio 91
Rudbeckia 90	Salsola 37	Sensitive Fern 2
Rue, Early Meadow 43	Saltwort 37	Service Berry 49
Family 56	Sambucus 83	Sessile-leaved Bell-
Meadow 43	Sandalwood Family 34	
Spleenwort 2	Sand-bar Willow 30	Setaria 16
Tall Meadow 43	Sandbur 11	Shad Bush 49
Rum Cherry 51	Sand Cherry 51	Shagbark Hickory 31
Rumex 36	Violet 62	Sharp-fruited Rush 24
Running Raspberry 52		Sharp-lobed Liver-
Swamp Blackberry 5		leaf 41
Rush 24	Rock 38	Sheep Laurel 69
Beak 22	Thyme-leaved 38	Sorrel 36
Bog 24	Sanguinaria 43	
Brown Beak 22	Sanguisorba 53	Sheepberry 83
Brownish-fruited 24	Sanicle 67	Sheep's Fescue 12
Chair-maker's 22	Sanicula 67	Shellbark Hickory 31
Club 22	Santalaceae 34	Shepherdia 64
		Shepherd's Purse 45
Common Scouring 4 Common Wood 24	Sapindaceae 55 Saponaria 39	Shield Fern 1
	Sarracenia 47	Water 40
Family 24		Shin Leaf 70
Grass 16	Sarraceniaceae 47	Shining Club Moss 5
Hairy Wood 24 Jointed 24	Sarsaparilla, Bristly 69 Wild 66	-01
		Willow 30
Knotted 24	Satureja 77	Shore Shad Bush 49
Scouring 5	Savin 6	Horsetail 5
Sharp-fruited 24	Savory 77	Short-stalked False
Slender 24	Summer 77	Pimpernel 79
Spike 21	Saxifraga 48	Short-styled Snake-
Toad 24	Saxifragaceae 48	root 67
White Beak 22	Saxifrage 48	Showy Lady's Slip-
Wood 24	Early 48	per 27
Russian Thistle 37	Family 48	Orchis 29
Rusty Woodsia 3	Golden 48	Tick Trefoil 53
Rutabaga 45	Swamp 48	Shrubby Bitter-sweet 58
Rutaceae 56	Scabious, Sweet 88	Cinquefoil 51
Rye 16	Scheuchzeria 9	Sicyos 84
Grass 13	Scirpus 22	Side-saddle Flower 47
Nodding Wild 12	Scoke, Common 37	Silene 39
Slender Wild 12	Scorpion-grass 75	Silkweed 73
Wild 12	Scouring Rush 5	Common 73
Rynchospora 22	Common 4	Silky Cornel 68
Sage, Flowering 76	Scrophularia 80	Dogwood 68
Wood 78	Schrophulariaceae 79	Willow 31
Sagittaria 9	Scutellaria 77	Silphium 91

Silver Maple 59	Water 35	Speedwell 80
Weed 50	Smilacina 26	Common 80
Silver-leaved Poplar 29	Smilax 26	Corn 80
Silvery Cinquefoil 50	Smooth Alder 32	Marsh 80
Spleenwort 2	Rock Cress 44	Purslane 80
Sinuate-leaved Even-	Rose 52	Thyme-leaved 80
ing Primrose 65	Sumach 58	Spergula 39
Sisymbrium 46	Yellow Violet 63	Sphenopholis 16
Sisymprium 45 Sisyrinchium 27	Smoother Sweet Cice-	Spice Bush 43
Sium 68	ly 67	Spike Rush 21
Skullcap 77	Snakehead 79	Spikenard 66
Mad-dog 77	Snake-mouth 29	False 26
Marsh 77		Spinulose Shield Fern 1
Small 77	Clustered 67	Spiny-leaved Sow
Skunk Cabbage 23	Large-fruited 67	Thistle 92
Cloopy Cotchfly 20	Seneca 57	Spiraea 53
Sleepy Catchfly 39 Slender Gerardia 79	Short-styled 67	Ash-leaved 53
Ladies' Tresses 29	White 88	Spiranthes 29
Mountain Rice 14	Snowberry 83	Spirodela 23
Muhlenbergia 14	Creeping 69	Spleenwort 2
Nettle 34	Garden 83	Ebony 2
Rock Brake 2	Soapherry Family 59	Maidenhair 2
Rush 24	Soapwort 39	Narrow-leaved 2
Vetch 55	Solanaceae 78	Rue 2
Wheat Grass 9	Solanum 78	Silvery 2
Wild Rye 12	Solidago 91	Sporobolus 16
Willow 30	Solomon's Seal 26	Spotted Cowbane 66
Wood Reed Grass 11		Dead Nettle 76
Slippery Elm 34	Small 26	Medick 54
Slough Grass 16	Star-flowered 26	Mint 76
Small Bedstraw 82	Three-leaved 26	St. John's-wort 62
Bitter Cress 45	Two-leaved 25	Touch-me-not 60
Crab Grass 12	Sonchus 92	Spreading Dogbane 72
Cranberry 71	Sorbaria 53	Spring Beauty 40
Red Morning Glory 73		Broad-leaved 40
Skullcap 77	Sorrel 36	Spruce 6
Solomon's Seal 26	Common Wood 56	Black 6
Sundrops 65	Field 36	Bog 6
White Aster 87	Garden 36	Cat 6
Whorled Pogonia 29	Lady's 56	Red 7
Wood Orchis 28	Sheep 36	White 6
Yellow Pond Lily 40	Wood 56	Spurge 57
Smaller Enchanter's	Yellow Wood 56	Cypress 57
Nightshade 64	Sour Gum 68	Family 57
Purple Fringed Or-	Sour-top Blueberry 70	Flowering 57
chis 28	Sow Thistle 92	Spurrey 39
Red-root 60	Sparganiaceae 7	Corn 39
Yellow Lady's Slip-	Sparganium 7	Field 39
per 27	Spartina 16	Squaw-weed 91
Small-flowered Crow-	Spatter Dock 40	Robbin's 91
foot 42	Spear Grass 15	Round-leaved 91
Evening Primrose 65	Spearmint 76	Squirrel Corn 44
Gerardia 79	Spearwort, Creeping 4	2Squirrel-tail Grass 13
St. John's-wort 62	Speckled Alder 31	Stachys 77
Smartweed, Common 3	5Specularia 84	Staff Tree 58

121

	221222	
Family 58	Staghorn 58	Blue Lettuce 90
Staghorn Sumach 58	Summer Grape 60	Buttercup 42
Staphylea 59	Savory 77	Cinquefoil 51
Staphyleaceae 59	Sundew 47	Cone-flower 90
Star Cucumber 84	Family 47	Crowfoot 42
Flower 72	Long-leaved 47	Green Orchis 28
Thistle 87	Round-leaved 47	Hairy Agrimony 49
Star-flowered Solomon'		Hairy Golden-rod 92
Seal 26	Sunflower 89	Meadow Rue 43
Starwort 39	Common 89	Oat Grass 10
Lesser 39	Pale-leaved Wood 89	White Aster 86
Water 57	Rough 89	White Lettuce 90
Steeple Bush 53	Thin-leaved 89	Wild Nettle 34
Steironema 71	Woodland 89	Wormwood 86
Stellaria 39	Swamp Beggar's-	Taller Fescue 12
Stemless Lady's Slip-	ticks 87	Tamarack 6
per 27	Blueberry 70	Tanacetum 92
Stickseed 74	Buttercup 43	Tansy 92
European 74	Dewberry 52	Common 92
Stick-tight 87	Dock 36	Tape Grass 9
Stiff Aster 86	Hickory 31	Taraxacum 92 Tare 55
Club Moss 5	Loosestrife 64	
Gentian 72 Water Crowfoot 42	Maple 59 Milkweed 73	Tartarian Honey- suckle 83
Stitchwort, Lesser 39	Persicaria 35	Taxaceae 6
Long-leaved 39	Pink 70	Taxus 6
Northern 39	Red Currant 48	Teaberry 69
Stone Clover 55	Rose 52	Tea, Labrador 69
Stonecrop 47	Saxifrage 48	Mexican 36
Ditch 47	White Oak 32	New Jersey 60
Mossy 47	Sweet Alyssum 46	Tear-thumb, Arrow-
Storksbill 56	Birch 32	leaved 36
Musk 56	Cicely 67	Halberd-leaved 35
Stout Ragged Golden-	Clover 54	Tendril Creeper 60
rod 92	Fern 31	Ternate Grape Fern 4
Stramonium 78	Flag 23	Teucrium 78
Strawberry 50	Gale 31	Thalictrum 43
Barren 53	Gale Family 31	Thimbleberry 52
Blite 37	Scabious 88	Thimble-weed 90
Tomato 78	Vernal Grass 10	Thin Grass 10
Wild 50	White Violet 62	Thin-leaved Sun-
Wood 50	William 38	flower 89 Thistle, Bull 88
Straw-colored Cy-	William Catchfly 39	Canada 88
perus 21	Sweetbrier 52 Sweet-scented Bed-	Common 88
Streptopus 26 Striped Maple 59	straw 82	Common Sow 92
Striped Maple 55 Strong-scented Era-	White Water Lily 40	Field Sow 92
grostis 12	Sycamore 49	Plumed 88
Succory 87	Symphoricarpos 83	Plumeless 87
Sugar Maple 59	Symphytum 75	Russian 37
Sugarberry 33	Symplocarpus 23	Sow 92
Sumach 58	Syringa 72	Spiny-leaved Sow 92
Dwarf 58	Tacamahac 29	Star 87
Poison 58	Taenidia 68	Thlaspi 47
Smooth 58	Tall Anemone 41	Thorn Apple 78

White 49	Trillium 26	glass 84
Thorny Amaranth 37	Nodding 26	Veratrum 26
Thoroughwort 88	Painted 26	Verbascum 80
Three-leaved Solomon's	Purple 26	Verbena 75
Seal 26	White 26	Verbenaceae 75
Three-seeded Mercury 5	7Triosteum 83	Veronica 80
Thuja 7	Triple-awned Grass 10	Vervain 75
Thyme, Basil 77	Trisetum 16	Blue 75
Thymelaeaceae 64	Triticum 16	Family 75
Thyme-leaved Sand-	True Forget-me-not 75	White 75
wort 38	Trumpet Weed 88	Vetch 55
Speedwell 80	Tsuga 7	Blue 55
Tiarella 48	Tuberous White Water	Common 55
Tick Trefoil 53	Lily 40	Milk 53
Tiger Lily 25	Tufted Loosestrife 71	Slender 55
Tilia 61	Tumble Mustard 46	Vetchling 54
Tiliaceae 61	Weed 37	Cream-colored 54
Timothy 15	Tupelo 68	Viburnum 83
Tinker's Weed 83	Turnip 45	Vicia 55
Toad Rush 24	Indian 23	Vinca 73
Toadflax 79	Wild 23	Vine, Allegheny 44
Bastard 34	Turtlehead 79	Common Matri-
Tobacco, Indian 84	Tussilago 92	mony 78
Ladies' 85	Twayblade 28, 29	Family 60
Tomatillo 78	Heart-leaved 29	Love 73
Tomato, Husk 78	Large 28	Matrimony 78
Strawberry 78	Twin-flower 82	Viola 62
Toothache Tree 56	Twisted Whitlow-	Violaceae 62
Toothed Arrow-wood 83		Violet 62
Toothwort 45	Twisted-stalk 26	Canada 62
Cut-leaved 45	Two-leaved Miter-	Dog 62
Large 45	wort 48	Dog's-tooth 25
Two-leaved 45	Pepper-root 45	Downy Yellow 63
Touch-me-not Family 6		Early Yellow 63
Pale 60	Toothwort 45	Family 62
Spotted 60	Typha 7	Lance-leaved 63
Tower Mustard 44	Typhaceae 7	Long-spurred 63
Tragopogon 92	Ulmus 34	Marsh Blue 62
Trailing Arbutus 69	Umbelliferae 66	Round-leaved 63
Traveler's Joy 41	Upright Bindweed 73	Sand 62
Treacle Mustard 46	Goosefoot 37	Smooth Yellow 63
Tree Club Moss 5	Urtica 34	Sweet White 62
Cranberry 83	Urticaceae 33	White 63
Nettle 33	Utricularia 80	Woolly Blue 63
Plane 49	Uvularia 26	Viper's Bugloss 74
Rowan 51	Vaccinium 70	Virginia Creeper 60
Staff 58	Valerian 84	Chain Fern 3
Toothache 56	Family 84	Virginian Cowslip 75
Trefoil 55	Valeriana 84	Virgin's Bower 41
Showy Tick 53	Valerianaceae 84	Vitaceae 60
Tick 53	Vallisneria 9	Vitis 60
Trembling Poplar 30	Vanilla Grass 13	Wake Robin 26
Trichostema 78	Velvet Leaf 61	Walking Loof 2
Trientalis 72	Velvet-leaf Blueberry 70	Wolnut 21
Trifolium 55	Venus's Looking-	Walnut 31

	11(1)1311	12
Family 31	Campion 38	Ginger 34
White 31	Cedar 7	Iris 27
Wartweed 57	Clover 55	Leek 25
Water Arum 23	Daisy 87	Lettuce 90
Avens 50	Elm 34	Lily of the Valley 25
Beech 32	Fringed Orchis 28	Liquorice 82
Cress 46	Golden-rod 91	Lupine 54
Hemlock 66	Grass 13	Madder 82
Hemp 37	Lettuce 90	Mint 76
Horehound 76	Maple 59	Morning Glory 73
Lily 40	Melilot 54	Oat Grass 11
Lily Family 40	Mulberry 33	Orange-red Lily 25
Marigold 87	Oak 32	Parsnip 67
Milfoil 65	Pine 7	Peppergrass 46
Milfoil Family 65	Poplar 29	Plum 51
Nymph 40	Snakeroot 88	Raisin 83
Oats 16	Spruce 6	Red Cherry 51
Parsnip 68	Sweet Clover 54	Red Raspberry 52
Pennywort 67	Thorn 49	Rice 16
Pepper 35	Trillium 26	Rye 12
Plantain 9	Vervain 75	Sarsaparilla 66
Plantain Family 9	Violet 63	Strawberry 50
Purslane 65	Walnut 31	Turnip 23
Rice 16	Willow 30	Yellow Lily 25
Shield 40	Wood Aster 86	Willow 30
Smartweed 35	White-grained Moun-	Beaked 30
Star Grass 24	tain Rice 14	Black 30
Starwort 57	White-heart Hickory 3	1 Bog 30
Starwort Family 57	Whiteweed 87	Crack 30
Willow 64	Whitlow-grass 46	Family 29
Waterleaf 74	Twisted 46	Glaucous 30
Family 74	Whorled Loosestrife 7	
Water-weed 9	Mallow 61	Prairie 30
Wavy-leaved Aster 87	Mint 76	Purple 30
Waxwork 58	Pogonia 29	Pussy 30
Weak Spear Grass 15	Wicky 69	Sand-bar 30
Weed, Bugle 75, 76	Wicopy 64	Shining 30
Carpet 38	Wild Allspice 43	Silky 30
Cut-leaved Bugle 76	Balsam-apple 84	Slender 30
Jamestown 78	Bean 53	Water 64
Jimson 78	Bergamot 76	White 30
Joe Pye 88	Black Cherry 51	Willow-herb 64
Silver 50	Black Currant 48	Great 65
Tinker's 83	Broom Corn 15	Northern 64
Trumpet 88	Calla 23	Purple-leaved 65
Tumble 37	Carrot 67	Wind Flower 41
Wheat 16	Chamomile 90	Winter Cress 44
Cow 79	Chess 11	Winterberry 58
White Ash 72	Clematis 41	Wintergreen 70
Avens 50	Coffee 83	Aromatic 69
Baneberry 40	Columbine 41	Bog 70
Beak Rush 22	Comfrey 74	Chickweed 72
Bent Grass 10	Cranesbill 56	Flowering 57
Birch 32	Cucumber 84	Greenish-flowered 70
Bog Orchis 28	Elder 66	One-sided 70

Round-leaved 70 Wire Grass 15 Witch Hobble 83 Witch-hazel 49 Family 49 Withe-rod 83 Wood Anemone 41 Betony 79 Chess 11 Fern 1 Grass 16 Horsetail 5 Lily 25 Nettle 33 Reed Grass 11 Rush 24 Sage 78 Sorrel 56 Sorrel Family 56 Strawberry 50 Woodbine 60 Woodland Agrimony 49 Sunflower 89 Woodsia 3

Blunt-lobed 3 Rustv 3 Woodwardia 3 Wool Grass 22 Woolly Blue Violet 63 Sweet Cicely 67 Worm-seed Mustard 46 Wormwood 86 Roman 85 Tall 86 Xanthium 93 Yarrow 85 Common 85 Yellow Adder's Tongue 25 Avens 50 Bartonia 72 Birch 32 Clintonia 25

Clover 55

Cress 46

Daisy 90

Dock 36

Pond Lily 40 Puccoon 42 Rocket 45 Sweet Clover 54 Water Crowfoot 42 Wood Sorrel 56 Yellow-barked Oak 33 Yew 6 American 6 Family 6 Zanthoxylum 56 Zea 16 Zig-zag Clover 55 Golden-rod 91 Zizania 16 Zizia 68

Foxtail 16

Melilot 54

Myrtle 71

Nut Grass 21

Pimpernel 68

Iris 27

Oak 33











North Carolina State University Libraries

QK190 .F6
FLORA OF BURLINGTON AND VICINITY A LIST OF TH